

表 2316 身長(cm) 透析前リン濃度(mg/dL)別 (透析患者全体)

透析前リン濃度	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
3未満	28	58	177	579	1,114	1,630	1,674	1,902	1,456	805	259	63	9	3	9,757	1,129	10,886	157.72	9.96
(%)	(0.3)	(0.6)	(1.8)	(5.9)	(11.4)	(16.7)	(17.2)	(19.5)	(14.9)	(8.3)	(2.7)	(0.6)	(0.1)	(0.0)	(100.0)				
3.0～	28	56	183	650	1,308	1,898	2,127	2,567	2,024	1,137	330	87	5	3	12,403	1,179	13,582	158.44	9.68
(%)	(0.2)	(0.5)	(1.5)	(5.2)	(10.5)	(15.3)	(17.1)	(20.7)	(16.3)	(9.2)	(2.7)	(0.7)	(0.0)	(0.0)	(100.0)				
3.5～	28	79	287	1,050	2,186	3,322	3,745	4,464	3,691	2,166	672	169	23	2	21,884	1,798	23,682	158.98	9.54
(%)	(0.1)	(0.4)	(1.3)	(4.8)	(10.0)	(15.2)	(17.1)	(20.4)	(16.9)	(9.9)	(3.1)	(0.8)	(0.1)	(0.0)	(100.0)				
4.0～	49	88	426	1,466	3,219	4,852	5,581	6,596	5,742	3,344	1,107	280	45	2	32,797	2,485	35,282	159.27	9.59
(%)	(0.1)	(0.3)	(1.3)	(4.5)	(9.8)	(14.8)	(17.0)	(20.1)	(17.5)	(10.2)	(3.4)	(0.9)	(0.1)	(0.0)	(100.0)				
4.5～	43	105	443	1,539	3,590	5,813	6,776	7,801	6,834	4,339	1,534	361	52	11	39,241	2,818	42,059	159.71	9.56
(%)	(0.1)	(0.3)	(1.1)	(3.9)	(9.1)	(14.8)	(17.3)	(19.9)	(17.4)	(11.1)	(3.9)	(0.9)	(0.1)	(0.0)	(100.0)				
5.0～	46	97	409	1,510	3,505	5,810	6,584	8,049	7,178	4,571	1,649	420	69	12	39,909	2,809	42,718	160.04	9.54
(%)	(0.1)	(0.2)	(1.0)	(3.8)	(8.8)	(14.6)	(16.5)	(20.2)	(18.0)	(11.5)	(4.1)	(1.1)	(0.2)	(0.0)	(100.0)				
5.5～	33	70	314	1,161	2,888	4,877	5,710	6,849	6,296	4,113	1,591	408	65	17	34,392	2,313	36,705	160.45	9.53
(%)	(0.1)	(0.2)	(0.9)	(3.4)	(8.4)	(14.2)	(16.6)	(19.9)	(18.3)	(12.0)	(4.6)	(1.2)	(0.2)	(0.0)	(100.0)				
6.0～	9	10	45	197	485	829	937	1,172	1,102	693	297	74	16	3	5,869	392	6,261	160.62	9.67
(%)	(0.2)	(0.2)	(0.8)	(3.4)	(8.3)	(14.1)	(16.0)	(20.0)	(18.8)	(11.8)	(5.1)	(1.3)	(0.3)	(0.1)	(100.0)				
6.1～	20	35	141	642	1,549	2,514	3,169	3,941	3,771	2,605	974	264	38	4	19,667	1,442	21,109	160.99	9.55
(%)	(0.1)	(0.2)	(0.7)	(3.3)	(7.9)	(12.8)	(16.1)	(20.0)	(19.2)	(13.2)	(5.0)	(1.3)	(0.2)	(0.0)	(100.0)				
6.5～	22	37	117	451	1,314	2,195	2,750	3,363	3,286	2,316	921	256	38	4	17,070	1,208	18,278	161.25	9.56
(%)	(0.1)	(0.2)	(0.7)	(2.6)	(7.7)	(12.9)	(16.1)	(19.7)	(19.3)	(13.6)	(5.4)	(1.5)	(0.2)	(0.0)	(100.0)				
7.0～	18	20	90	321	731	1,280	1,655	2,107	1,984	1,525	570	168	27	2	10,498	779	11,277	161.44	9.65
(%)	(0.2)	(0.2)	(0.9)	(3.1)	(7.0)	(12.2)	(15.8)	(20.1)	(18.9)	(14.5)	(5.4)	(1.6)	(0.3)	(0.0)	(100.0)				
7.5～	8	8	59	177	444	765	999	1,260	1,248	941	387	116	18	3	6,433	425	6,858	161.75	9.85
(%)	(0.1)	(0.1)	(0.9)	(2.8)	(6.9)	(11.9)	(15.5)	(19.6)	(19.4)	(14.6)	(6.0)	(1.8)	(0.3)	(0.0)	(100.0)				
8.0～	4	5	29	113	244	458	536	724	729	597	245	76	19	4	3,783	270	4,053	162.19	9.81
(%)	(0.1)	(0.1)	(0.8)	(3.0)	(6.4)	(12.1)	(14.2)	(19.1)	(19.3)	(15.8)	(6.5)	(2.0)	(0.5)	(0.1)	(100.0)				
8.5～	3	4	11	56	151	287	324	464	486	356	163	56	11	4	2,376	196	2,572	162.47	9.75
(%)	(0.1)	(0.2)	(0.5)	(2.4)	(6.4)	(12.1)	(13.6)	(19.5)	(20.5)	(15.0)	(6.9)	(2.4)	(0.5)	(0.2)	(100.0)				
9.0～	7	8	27	92	222	437	528	709	747	626	273	99	17	1	3,793	284	4,077	162.65	9.82
(%)	(0.2)	(0.2)	(0.7)	(2.4)	(5.9)	(11.5)	(13.9)	(18.7)	(19.7)	(16.5)	(7.2)	(2.6)	(0.4)	(0.0)	(100.0)				
合計	346	680	2,758	10,004	22,950	36,967	43,095	51,968	46,574	30,134	10,972	2,897	452	75	259,872	19,527	279,399	160.05	9.65
(%)	(0.1)	(0.3)	(1.1)	(3.8)	(8.8)	(14.2)	(16.6)	(20.0)	(17.9)	(11.6)	(4.2)	(1.1)	(0.2)	(0.0)	(100.0)				
記載なし	8	16	79	293	723	1,268	1,377	1,682	1,414	940	347	82	13	2	8,244	22,465	30,709	159.90	9.59
(%)	(0.1)	(0.2)	(1.0)	(3.6)	(8.8)	(15.4)	(16.7)	(20.4)	(17.2)	(11.4)	(4.2)	(1.0)	(0.2)	(0.0)	(100.0)				
総計	354	696	2,837	10,297	23,673	38,235	44,472	53,650	47,988	31,074	11,319	2,979	465	77	268,116	41,992	310,108	160.05	9.65
(%)	(0.1)	(0.3)	(1.1)	(3.8)	(8.8)	(14.3)	(16.6)	(20.0)	(17.9)	(11.6)	(4.2)	(1.1)	(0.2)	(0.0)	(100.0)				
平均	5.01	4.80	4.90	4.94	5.04	5.11	5.17	5.21	5.30	5.43	5.56	5.69	5.83	5.68	5.22	5.10	5.21		
標準偏差	1.63	1.45	1.41	1.39	1.37	1.39	1.39	1.40	1.42	1.46	1.49	1.57	1.58	1.64	1.42	1.50	1.43		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査