

表 2319 身長(cm) 透析前ヘモグロビン濃度(g/dL)別 (透析患者全体)

透析前ヘモグロビン濃度	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
5未満		2	2	6	29	49	40	64	48	27	13	2			282	21	303	159.55	9.34
(%)		(0.7)	(0.7)	(2.1)	(10.3)	(17.4)	(14.2)	(22.7)	(17.0)	(9.6)	(4.6)	(0.7)			(100.0)				
5～	1		3	20	30	73	69	73	60	40	16	1	1		387	42	429	159.15	9.28
(%)	(0.3)		(0.8)	(5.2)	(7.8)	(18.9)	(17.8)	(18.9)	(15.5)	(10.3)	(4.1)	(0.3)	(0.3)		(100.0)				
6～	2	2	17	55	112	156	179	194	183	91	38	9	2		1,040	120	1,160	158.65	10.09
(%)	(0.2)	(0.2)	(1.6)	(5.3)	(10.8)	(15.0)	(17.2)	(18.7)	(17.6)	(8.8)	(3.7)	(0.9)	(0.2)		(100.0)				
7～	11	20	57	191	443	648	685	772	602	393	131	33	9	1	3,996	439	4,435	158.51	10.03
(%)	(0.3)	(0.5)	(1.4)	(4.8)	(11.1)	(16.2)	(17.1)	(19.3)	(15.1)	(9.8)	(3.3)	(0.8)	(0.2)	(0.0)	(100.0)				
8～	27	68	230	702	1,421	2,096	2,247	2,572	2,104	1,285	446	112	18	1	13,329	1,225	14,554	158.52	9.94
(%)	(0.2)	(0.5)	(1.7)	(5.3)	(10.7)	(15.7)	(16.9)	(19.3)	(15.8)	(9.6)	(3.3)	(0.8)	(0.1)	(0.0)	(100.0)				
9～	65	116	526	1,930	4,370	6,803	7,436	8,258	7,097	4,289	1,423	349	56	16	42,734	3,255	45,989	159.01	9.62
(%)	(0.2)	(0.3)	(1.2)	(4.5)	(10.2)	(15.9)	(17.4)	(19.3)	(16.6)	(10.0)	(3.3)	(0.8)	(0.1)	(0.0)	(100.0)				
10～	111	209	926	3,420	7,952	12,867	14,760	17,612	15,510	9,896	3,542	931	141	22	87,899	6,013	93,912	159.88	9.61
(%)	(0.1)	(0.2)	(1.1)	(3.9)	(9.0)	(14.6)	(16.8)	(20.0)	(17.6)	(11.3)	(4.0)	(1.1)	(0.2)	(0.0)	(100.0)				
11～	84	181	669	2,487	5,973	9,885	11,881	14,911	13,537	8,998	3,470	901	140	23	73,140	5,033	78,173	160.60	9.58
(%)	(0.1)	(0.2)	(0.9)	(3.4)	(8.2)	(13.5)	(16.2)	(20.4)	(18.5)	(12.3)	(4.7)	(1.2)	(0.2)	(0.0)	(100.0)				
12～	28	51	253	849	1,957	3,241	4,311	5,486	5,304	3,512	1,281	385	60	8	26,726	2,128	28,854	161.13	9.47
(%)	(0.1)	(0.2)	(0.9)	(3.2)	(7.3)	(12.1)	(16.1)	(20.5)	(19.8)	(13.1)	(4.8)	(1.4)	(0.2)	(0.0)	(100.0)				
13～	6	17	57	215	448	726	959	1,354	1,405	1,004	376	110	13	1	6,691	617	7,308	161.80	9.60
(%)	(0.1)	(0.3)	(0.9)	(3.2)	(6.7)	(10.9)	(14.3)	(20.2)	(21.0)	(15.0)	(5.6)	(1.6)	(0.2)	(0.0)	(100.0)				
14～	8	7	15	63	122	235	352	472	518	440	179	56	9	3	2,479	194	2,673	162.99	10.03
(%)	(0.3)	(0.3)	(0.6)	(2.5)	(4.9)	(9.5)	(14.2)	(19.0)	(20.9)	(17.7)	(7.2)	(2.3)	(0.4)	(0.1)	(100.0)				
合計	343	673	2,755	9,938	22,857	36,779	42,919	51,768	46,368	29,975	10,915	2,889	449	75	258,703	19,087	277,790	160.05	9.65
(%)	(0.1)	(0.3)	(1.1)	(3.8)	(8.8)	(14.2)	(16.6)	(20.0)	(17.9)	(11.6)	(4.2)	(1.1)	(0.2)	(0.0)	(100.0)				
記載なし	11	23	82	359	816	1,456	1,553	1,882	1,620	1,099	404	90	16	2	9,413	22,905	32,318	159.91	9.64
(%)	(0.1)	(0.2)	(0.9)	(3.8)	(8.7)	(15.5)	(16.5)	(20.0)	(17.2)	(11.7)	(4.3)	(1.0)	(0.2)	(0.0)	(100.0)				
総計	354	696	2,837	10,297	23,673	38,235	44,472	53,650	47,988	31,074	11,319	2,979	465	77	268,116	41,992	310,108	160.05	9.65
(%)	(0.1)	(0.3)	(1.1)	(3.8)	(8.8)	(14.3)	(16.6)	(20.0)	(17.9)	(11.6)	(4.2)	(1.1)	(0.2)	(0.0)	(100.0)				
平均	10.51	10.51	10.50	10.54	10.56	10.59	10.67	10.73	10.80	10.86	10.90	10.98	10.92	10.88	10.71	10.67	10.71		
標準偏差	1.39	1.40	1.30	1.27	1.26	1.25	1.26	1.27	1.28	1.29	1.31	1.30	1.33	1.31	1.28	1.39	1.28		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査