

表 2322 身長(cm) Non-HDL-C濃度(mg/dL)別（透析患者全体）

Non-HDL-C濃度	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
50未満	4	3	21	72	179	328	506	740	700	416	173	36	7	1	3,186	224	3,410	162.00	8.85
(%)	(0.1)	(0.1)	(0.7)	(2.3)	(5.6)	(10.3)	(15.9)	(23.2)	(22.0)	(13.1)	(5.4)	(1.1)	(0.2)	(0.0)	(100.0)				
50～	16	30	103	387	941	1,592	2,319	3,508	3,313	1,993	727	214	25	7	15,175	926	16,101	161.77	9.11
(%)	(0.1)	(0.2)	(0.7)	(2.6)	(6.2)	(10.5)	(15.3)	(23.1)	(21.8)	(13.1)	(4.8)	(1.4)	(0.2)	(0.0)	(100.0)				
70～	45	83	338	1,143	2,443	4,124	5,434	7,179	6,663	4,027	1,504	386	52	9	33,430	2,139	35,569	160.80	9.45
(%)	(0.1)	(0.2)	(1.0)	(3.4)	(7.3)	(12.3)	(16.3)	(21.5)	(19.9)	(12.0)	(4.5)	(1.2)	(0.2)	(0.0)	(100.0)				
90～	43	103	467	1,581	3,460	5,506	6,516	8,107	7,334	4,677	1,655	435	65	12	39,961	2,417	42,378	160.14	9.57
(%)	(0.1)	(0.3)	(1.2)	(4.0)	(8.7)	(13.8)	(16.3)	(20.3)	(18.4)	(11.7)	(4.1)	(1.1)	(0.2)	(0.0)	(100.0)				
110～	38	81	367	1,372	3,215	4,938	5,343	6,241	5,339	3,603	1,241	355	68	9	32,210	1,993	34,203	159.53	9.74
(%)	(0.1)	(0.3)	(1.1)	(4.3)	(10.0)	(15.3)	(16.6)	(19.4)	(16.6)	(11.2)	(3.9)	(1.1)	(0.2)	(0.0)	(100.0)				
130～	17	54	236	855	2,077	3,234	3,371	3,581	3,111	2,161	786	220	34	3	19,740	1,223	20,963	159.25	9.75
(%)	(0.1)	(0.3)	(1.2)	(4.3)	(10.5)	(16.4)	(17.1)	(18.1)	(15.8)	(10.9)	(4.0)	(1.1)	(0.2)	(0.0)	(100.0)				
150～	15	29	117	475	1,082	1,698	1,709	1,736	1,502	1,041	415	113	21	2	9,955	635	10,590	158.98	9.95
(%)	(0.2)	(0.3)	(1.2)	(4.8)	(10.9)	(17.1)	(17.2)	(17.4)	(15.1)	(10.5)	(4.2)	(1.1)	(0.2)	(0.0)	(100.0)				
170～	6	11	46	198	502	800	767	725	652	453	184	51	11	1	4,407	315	4,722	158.80	10.00
(%)	(0.1)	(0.2)	(1.0)	(4.5)	(11.4)	(18.2)	(17.4)	(16.5)	(14.8)	(10.3)	(4.2)	(1.2)	(0.2)	(0.0)	(100.0)				
190～	5	10	15	75	183	294	323	290	226	172	60	20	3		1,676	133	1,809	158.61	9.90
(%)	(0.3)	(0.6)	(0.9)	(4.5)	(10.9)	(17.5)	(19.3)	(17.3)	(13.5)	(10.3)	(3.6)	(1.2)	(0.2)		(100.0)				
210～	2	4	13	46	118	195	177	172	128	112	43	8			1,018	106	1,124	158.37	10.15
(%)	(0.2)	(0.4)	(1.3)	(4.5)	(11.6)	(19.2)	(17.4)	(16.9)	(12.6)	(11.0)	(4.2)	(0.8)			(100.0)				
合計	191	408	1,723	6,204	14,200	22,709	26,465	32,279	28,968	18,655	6,788	1,838	286	44	160,758	10,111	170,869	160.10	9.63
(%)	(0.1)	(0.3)	(1.1)	(3.9)	(8.8)	(14.1)	(16.5)	(20.1)	(18.0)	(11.6)	(4.2)	(1.1)	(0.2)	(0.0)	(100.0)				
記載なし	163	288	1,114	4,093	9,473	15,526	18,007	21,371	19,020	12,419	4,531	1,141	179	33	107,358	31,881	139,239	159.96	9.69
(%)	(0.2)	(0.3)	(1.0)	(3.8)	(8.8)	(14.5)	(16.8)	(19.9)	(17.7)	(11.6)	(4.2)	(1.1)	(0.2)	(0.0)	(100.0)				
総計	354	696	2,837	10,297	23,673	38,235	44,472	53,650	47,988	31,074	11,319	2,979	465	77	268,116	41,992	310,108	160.05	9.65
(%)	(0.1)	(0.3)	(1.1)	(3.8)	(8.8)	(14.3)	(16.6)	(20.0)	(17.9)	(11.6)	(4.2)	(1.1)	(0.2)	(0.0)	(100.0)				
平均	109.99	112.05	110.42	111.96	112.73	111.88	108.05	104.19	103.12	105.01	105.02	105.91	109.73	98.82	107.02	108.00	107.08		
標準偏差	37.69	35.63	32.01	32.94	33.69	33.99	33.72	32.79	32.64	33.22	34.04	34.14	32.98	30.20	33.48	35.48	33.61		

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査