

表 2323 身長(cm) 透析前収縮期血圧(mmHg)別（透析患者全体）

透析前収縮期血圧	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
60未満	1		4	3	1	9	12	4	1	2	1				38	2	40	153.00	11.31
(%)	(2.6)		(10.5)	(7.9)	(2.6)	(23.7)	(31.6)	(10.5)	(2.6)	(5.3)	(2.6)				(100.0)				
60～	1	2	6	30	43	69	76	73	57	37	10	2			406	27	433	157.54	10.25
(%)	(0.2)	(0.5)	(1.5)	(7.4)	(10.6)	(17.0)	(18.7)	(18.0)	(14.0)	(9.1)	(2.5)	(0.5)			(100.0)				
80～	22	16	59	164	341	527	572	570	452	247	90	25	2	2	3,089	249	3,338	157.57	10.81
(%)	(0.7)	(0.5)	(1.9)	(5.3)	(11.0)	(17.1)	(18.5)	(18.5)	(14.6)	(8.0)	(2.9)	(0.8)	(0.1)	(0.1)	(100.0)				
100～	48	61	238	834	1,695	2,584	2,876	3,171	2,617	1,560	534	119	21	1	16,359	1,248	17,607	158.59	10.00
(%)	(0.3)	(0.4)	(1.5)	(5.1)	(10.4)	(15.8)	(17.6)	(19.4)	(16.0)	(9.5)	(3.3)	(0.7)	(0.1)	(0.0)	(100.0)				
120～	60	150	587	2,047	4,566	7,242	8,583	9,926	8,551	5,402	1,916	505	79	9	49,623	3,648	53,271	159.64	9.65
(%)	(0.1)	(0.3)	(1.2)	(4.1)	(9.2)	(14.6)	(17.3)	(20.0)	(17.2)	(10.9)	(3.9)	(1.0)	(0.2)	(0.0)	(100.0)				
140～	86	183	745	2,713	6,483	10,575	12,486	15,428	13,810	8,846	3,127	850	121	15	75,468	5,354	80,822	160.22	9.50
(%)	(0.1)	(0.2)	(1.0)	(3.6)	(8.6)	(14.0)	(16.5)	(20.4)	(18.3)	(11.7)	(4.1)	(1.1)	(0.2)	(0.0)	(100.0)				
160～	59	109	532	1,961	4,512	7,392	8,784	10,896	10,174	6,685	2,493	661	105	16	54,379	3,816	58,195	160.51	9.56
(%)	(0.1)	(0.2)	(1.0)	(3.6)	(8.3)	(13.6)	(16.2)	(20.0)	(18.7)	(12.3)	(4.6)	(1.2)	(0.2)	(0.0)	(100.0)				
180～	25	53	215	787	1,803	2,859	3,189	4,010	3,816	2,548	1,038	277	39	9	20,668	1,556	22,224	160.49	9.80
(%)	(0.1)	(0.3)	(1.0)	(3.8)	(8.7)	(13.8)	(15.4)	(19.4)	(18.5)	(12.3)	(5.0)	(1.3)	(0.2)	(0.0)	(100.0)				
200～	5	9	45	194	500	714	774	981	916	670	288	75	14	3	5,188	358	5,546	160.60	9.94
(%)	(0.1)	(0.2)	(0.9)	(3.7)	(9.6)	(13.8)	(14.9)	(18.9)	(17.7)	(12.9)	(5.6)	(1.4)	(0.3)	(0.1)	(100.0)				
220～	1	6	10	54	102	155	147	199	175	133	76	18	9		1,085	75	1,160	160.42	10.87
(%)	(0.1)	(0.6)	(0.9)	(5.0)	(9.4)	(14.3)	(13.5)	(18.3)	(16.1)	(12.3)	(7.0)	(1.7)	(0.8)		(100.0)				
合計	308	589	2,441	8,787	20,046	32,126	37,499	45,258	40,569	26,130	9,573	2,532	390	55	226,303	16,333	242,636	160.04	9.67
(%)	(0.1)	(0.3)	(1.1)	(3.9)	(8.9)	(14.2)	(16.6)	(20.0)	(17.9)	(11.5)	(4.2)	(1.1)	(0.2)	(0.0)	(100.0)				
記載なし	46	107	396	1,510	3,627	6,109	6,973	8,392	7,419	4,944	1,746	447	75	22	41,813	25,659	67,472	160.08	9.56
(%)	(0.1)	(0.3)	(0.9)	(3.6)	(8.7)	(14.6)	(16.7)	(20.1)	(17.7)	(11.8)	(4.2)	(1.1)	(0.2)	(0.1)	(100.0)				
総計	354	696	2,837	10,297	23,673	38,235	44,472	53,650	47,988	31,074	11,319	2,979	465	77	268,116	41,992	310,108	160.05	9.65
(%)	(0.1)	(0.3)	(1.1)	(3.8)	(8.8)	(14.3)	(16.6)	(20.0)	(17.9)	(11.6)	(4.2)	(1.1)	(0.2)	(0.0)	(100.0)				
平均	142.46	147.41	147.96	149.18	150.10	150.21	150.07	151.08	152.07	152.91	154.23	154.82	155.95	157.80	151.15	150.85	151.13		
標準偏差	29.81	25.95	25.95	25.68	25.08	24.69	24.13	23.78	23.55	23.67	24.18	23.74	25.36	25.93	24.21	24.51	24.23		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査