

表 2331 身長(cm) 透析後体重(kg)別（施設血液透析患者 透析回数週3回）

透析後体重	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
20未満	1		1	3	6	14	16	10	17	4	1	1			74	10	84	156.97	14.65
(%)	(1.4)		(1.4)	(4.1)	(8.1)	(18.9)	(21.6)	(13.5)	(23.0)	(5.4)	(1.4)	(1.4)			(100.0)				
20～	24	65	106	221	152	87	44	7	4	2	1	1			714	88	802	143.12	8.44
(%)	(3.4)	(9.1)	(14.8)	(31.0)	(21.3)	(12.2)	(6.2)	(1.0)	(0.6)	(0.3)	(0.1)	(0.1)			(100.0)				
30～	93	320	1,159	3,240	5,029	4,466	2,110	744	194	60	11	1	1	1	17,429	1,688	19,117	148.21	7.15
(%)	(0.5)	(1.8)	(6.6)	(18.6)	(28.9)	(25.6)	(12.1)	(4.3)	(1.1)	(0.3)	(0.1)	(0.0)	(0.0)	(0.0)	(100.0)				
40～	69	134	805	3,550	8,867	13,782	12,980	9,340	4,349	1,245	186	19	1	2	55,329	4,480	59,809	154.65	7.67
(%)	(0.1)	(0.2)	(1.5)	(6.4)	(16.0)	(24.9)	(23.5)	(16.9)	(7.9)	(2.3)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
50～	27	33	163	907	3,489	8,267	13,379	18,707	14,217	6,420	1,413	191	13	4	67,230	4,879	72,109	160.88	7.47
(%)	(0.0)	(0.0)	(0.2)	(1.3)	(5.2)	(12.3)	(19.9)	(27.8)	(21.1)	(9.5)	(2.1)	(0.3)	(0.0)	(0.0)	(100.0)				
60～	10	6	30	212	803	2,286	4,498	9,446	12,010	8,665	2,761	549	60	6	41,342	2,740	44,082	165.19	7.30
(%)	(0.0)	(0.0)	(0.1)	(0.5)	(1.9)	(5.5)	(10.9)	(22.8)	(29.1)	(21.0)	(6.7)	(1.3)	(0.1)	(0.0)	(100.0)				
70～	8	2	2	33	147	550	986	2,592	4,443	4,482	2,204	611	88	7	16,155	1,029	17,184	168.10	7.32
(%)	(0.0)	(0.0)	(0.0)	(0.2)	(0.9)	(3.4)	(6.1)	(16.0)	(27.5)	(27.7)	(13.6)	(3.8)	(0.5)	(0.0)	(100.0)				
80～	1			8	44	122	270	637	1,303	1,714	1,109	418	80	7	5,713	368	6,081	170.19	7.31
(%)	(0.0)			(0.1)	(0.8)	(2.1)	(4.7)	(11.2)	(22.8)	(30.0)	(19.4)	(7.3)	(1.4)	(0.1)	(100.0)				
90～	1		1	3	4	31	75	159	404	609	466	242	47	7	2,049	146	2,195	171.88	7.27
(%)	(0.0)		(0.0)	(0.1)	(0.2)	(1.5)	(3.7)	(7.8)	(19.7)	(29.7)	(22.7)	(11.8)	(2.3)	(0.3)	(100.0)				
100～				2	1	5	16	50	97	197	180	105	37	12	702	53	755	173.80	7.64
(%)				(0.3)	(0.1)	(0.7)	(2.3)	(7.1)	(13.8)	(28.1)	(25.6)	(15.0)	(5.3)	(1.7)	(100.0)				
110～					1	5	11	11	34	76	70	55	9	7	279	20	299	174.08	7.81
(%)					(0.4)	(1.8)	(3.9)	(3.9)	(12.2)	(27.2)	(25.1)	(19.7)	(3.2)	(2.5)	(100.0)				
120～			1				2	8	24	43	47	38	11	3	177	12	189	175.07	7.57
(%)			(0.6)				(1.1)	(4.5)	(13.6)	(24.3)	(26.6)	(21.5)	(6.2)	(1.7)	(100.0)				
合計	234	560	2,268	8,179	18,543	29,615	34,387	41,711	37,096	23,517	8,449	2,231	347	56	207,193	15,513	222,706	159.95	9.59
(%)	(0.1)	(0.3)	(1.1)	(3.9)	(8.9)	(14.3)	(16.6)	(20.1)	(17.9)	(11.4)	(4.1)	(1.1)	(0.2)	(0.0)	(100.0)				
記載なし	3	8	37	104	276	399	470	547	480	310	100	23	4	1	2,762	3,503	6,265	159.45	9.56
(%)	(0.1)	(0.3)	(1.3)	(3.8)	(10.0)	(14.4)	(17.0)	(19.8)	(17.4)	(11.2)	(3.6)	(0.8)	(0.1)	(0.0)	(100.0)				
総計	237	568	2,305	8,283	18,819	30,014	34,857	42,258	37,576	23,827	8,549	2,254	351	57	209,955	19,016	228,971	159.94	9.59
(%)	(0.1)	(0.3)	(1.1)	(3.9)	(9.0)	(14.3)	(16.6)	(20.1)	(17.9)	(11.3)	(4.1)	(1.1)	(0.2)	(0.0)	(100.0)				
平均	41.85	37.69	40.01	42.29	45.13	48.46	52.11	56.52	60.98	65.83	71.29	77.84	83.19	88.89	55.55	54.08	55.45		
標準偏差	12.46	7.60	8.84	7.97	8.38	9.07	9.28	9.58	10.63	12.00	13.70	15.80	16.69	23.73	12.89	12.97	12.90		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査