

表 2332 身長(cm) 透析後体重(kg)別（施設血液透析患者 男性のみ 透析回数週3回）

透析後体重	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
20未満	1			1		1	8	10	16	4	1	1			43	6	49	161.21	17.48
(%)	(2.3)			(2.3)		(2.3)	(18.6)	(23.3)	(37.2)	(9.3)	(2.3)	(2.3)			(100.0)				
20～	4	3	2	6	4	3	3	1	4	2	1	1			34	6	40	148.38	16.95
(%)	(11.8)	(8.8)	(5.9)	(17.6)	(11.8)	(8.8)	(8.8)	(2.9)	(11.8)	(5.9)	(2.9)	(2.9)			(100.0)				
30～	18	12	38	117	298	618	661	467	154	53	7	1	1	1	2,446	272	2,718	154.78	8.07
(%)	(0.7)	(0.5)	(1.6)	(4.8)	(12.2)	(25.3)	(27.0)	(19.1)	(6.3)	(2.2)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
40～	33	12	24	183	870	3,357	6,577	7,321	4,029	1,206	178	18	1	2	23,811	2,100	25,911	159.67	6.55
(%)	(0.1)	(0.1)	(0.1)	(0.8)	(3.7)	(14.1)	(27.6)	(30.7)	(16.9)	(5.1)	(0.7)	(0.1)	(0.0)	(0.0)	(100.0)				
50～	15	13	20	55	360	2,554	8,482	16,366	13,667	6,337	1,399	190	13	4	49,475	3,652	53,127	163.48	6.07
(%)	(0.0)	(0.0)	(0.0)	(0.1)	(0.7)	(5.2)	(17.1)	(33.1)	(27.6)	(12.8)	(2.8)	(0.4)	(0.0)	(0.0)	(100.0)				
60～	6	3	9	21	63	506	2,652	8,382	11,630	8,571	2,742	547	60	6	35,198	2,318	37,516	166.90	5.96
(%)	(0.0)	(0.0)	(0.0)	(0.1)	(0.2)	(1.4)	(7.5)	(23.8)	(33.0)	(24.4)	(7.8)	(1.6)	(0.2)	(0.0)	(100.0)				
70～	8	1		3	14	62	434	2,138	4,272	4,436	2,186	608	86	7	14,255	892	15,147	169.54	6.11
(%)	(0.1)	(0.0)		(0.0)	(0.1)	(0.4)	(3.0)	(15.0)	(30.0)	(31.1)	(15.3)	(4.3)	(0.6)	(0.0)	(100.0)				
80～	1			2	3	11	70	470	1,229	1,695	1,104	417	80	7	5,089	329	5,418	171.63	6.04
(%)	(0.0)			(0.0)	(0.1)	(0.2)	(1.4)	(9.2)	(24.2)	(33.3)	(21.7)	(8.2)	(1.6)	(0.1)	(100.0)				
90～				2		2	23	95	368	600	464	242	46	7	1,849	125	1,974	173.13	6.12
(%)				(0.1)		(0.1)	(1.2)	(5.1)	(19.9)	(32.4)	(25.1)	(13.1)	(2.5)	(0.4)	(100.0)				
100～						2	1	29	89	192	179	105	37	12	646	48	694	174.93	6.61
(%)						(0.3)	(0.2)	(4.5)	(13.8)	(29.7)	(27.7)	(16.3)	(5.7)	(1.9)	(100.0)				
110～							3	5	33	76	69	55	9	7	257	16	273	175.41	6.35
(%)							(1.2)	(1.9)	(12.8)	(29.6)	(26.8)	(21.4)	(3.5)	(2.7)	(100.0)				
120～			1					4	21	43	46	38	11	3	167	11	178	175.69	7.19
(%)			(0.6)					(2.4)	(12.6)	(25.7)	(27.5)	(22.8)	(6.6)	(1.8)	(100.0)				
合計	86	44	94	390	1,612	7,116	18,914	35,288	35,512	23,215	8,376	2,223	344	56	133,270	9,775	143,045	164.72	7.26
(%)	(0.1)	(0.0)	(0.1)	(0.3)	(1.2)	(5.3)	(14.2)	(26.5)	(26.6)	(17.4)	(6.3)	(1.7)	(0.3)	(0.0)	(100.0)				
記載なし	1	1		7	22	97	248	456	459	305	99	23	4		1,722	2,266	3,988	164.42	7.11
(%)	(0.1)	(0.1)		(0.4)	(1.3)	(5.6)	(14.4)	(26.5)	(26.7)	(17.7)	(5.7)	(1.3)	(0.2)		(100.0)				
総計	87	45	94	397	1,634	7,213	19,162	35,744	35,971	23,520	8,475	2,246	348	56	134,992	12,041	147,033	164.72	7.26
(%)	(0.1)	(0.0)	(0.1)	(0.3)	(1.2)	(5.3)	(14.2)	(26.5)	(26.6)	(17.4)	(6.3)	(1.7)	(0.3)	(0.0)	(100.0)				
平均	47.78	46.00	47.14	44.55	46.27	49.39	52.73	56.72	61.03	65.86	71.34	77.88	83.20	88.89	59.68	58.27	59.58		
標準偏差	14.16	10.76	26.22	9.70	7.61	7.52	8.13	9.09	10.51	11.98	13.65	15.80	16.76	23.73	12.17	12.24	12.18		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査