

表 2340 身長(cm) Body Mass Index別（施設血液透析患者 透析回数週3回）

Body Mass Index	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
14未満	1	3	18	77	178	312	278	187	128	83	32	7	2	5	1,311		1,311	156.41	9.23
(%)	(0.1)	(0.2)	(1.4)	(5.9)	(13.6)	(23.8)	(21.2)	(14.3)	(9.8)	(6.3)	(2.4)	(0.5)	(0.2)	(0.4)	(100.0)				
14～	6	28	106	479	1,081	1,578	1,491	1,207	937	505	143	38	5	2	7,606		7,606	156.51	9.15
(%)	(0.1)	(0.4)	(1.4)	(6.3)	(14.2)	(20.7)	(19.6)	(15.9)	(12.3)	(6.6)	(1.9)	(0.5)	(0.1)	(0.0)	(100.0)				
16～	8	84	321	1,252	2,894	4,523	4,547	4,539	3,547	1,950	662	135	15	3	24,480		24,480	157.83	9.22
(%)	(0.0)	(0.3)	(1.3)	(5.1)	(11.8)	(18.5)	(18.6)	(18.5)	(14.5)	(8.0)	(2.7)	(0.6)	(0.1)	(0.0)	(100.0)				
18～	27	117	521	1,852	4,414	6,708	7,686	8,690	7,290	4,137	1,353	308	49	4	43,156		43,156	159.05	9.28
(%)	(0.1)	(0.3)	(1.2)	(4.3)	(10.2)	(15.5)	(17.8)	(20.1)	(16.9)	(9.6)	(3.1)	(0.7)	(0.1)	(0.0)	(100.0)				
20～	28	116	486	1,803	3,894	6,264	8,158	10,382	8,803	5,494	1,780	410	55	7	47,680		47,680	160.20	9.20
(%)	(0.1)	(0.2)	(1.0)	(3.8)	(8.2)	(13.1)	(17.1)	(21.8)	(18.5)	(11.5)	(3.7)	(0.9)	(0.1)	(0.0)	(100.0)				
22～	20	90	366	1,244	2,750	4,728	5,714	7,850	7,293	4,688	1,595	403	65	4	36,810		36,810	160.84	9.30
(%)	(0.1)	(0.2)	(1.0)	(3.4)	(7.5)	(12.8)	(15.5)	(21.3)	(19.8)	(12.7)	(4.3)	(1.1)	(0.2)	(0.0)	(100.0)				
24～	15	43	217	698	1,603	2,551	3,341	4,509	4,414	2,962	1,175	324	53	8	21,913		21,913	161.40	9.51
(%)	(0.1)	(0.2)	(1.0)	(3.2)	(7.3)	(11.6)	(15.2)	(20.6)	(20.1)	(13.5)	(5.4)	(1.5)	(0.2)	(0.0)	(100.0)				
26～	16	32	106	372	836	1,414	1,596	2,234	2,372	1,698	736	233	39	4	11,688		11,688	161.80	9.83
(%)	(0.1)	(0.3)	(0.9)	(3.2)	(7.2)	(12.1)	(13.7)	(19.1)	(20.3)	(14.5)	(6.3)	(2.0)	(0.3)	(0.0)	(100.0)				
28～	15	20	63	169	439	761	757	1,082	1,111	939	445	153	27	10	5,991		5,991	162.11	10.38
(%)	(0.3)	(0.3)	(1.1)	(2.8)	(7.3)	(12.7)	(12.6)	(18.1)	(18.5)	(15.7)	(7.4)	(2.6)	(0.5)	(0.2)	(100.0)				
30～	25	19	53	187	370	624	641	839	975	827	405	164	28	7	5,164		5,164	162.15	10.97
(%)	(0.5)	(0.4)	(1.0)	(3.6)	(7.2)	(12.1)	(12.4)	(16.2)	(18.9)	(16.0)	(7.8)	(3.2)	(0.5)	(0.1)	(100.0)				
合計	161	552	2,257	8,133	18,459	29,463	34,209	41,519	36,870	23,283	8,326	2,175	338	54	205,799		205,799	159.95	9.49
(%)	(0.1)	(0.3)	(1.1)	(4.0)	(9.0)	(14.3)	(16.6)	(20.2)	(17.9)	(11.3)	(4.0)	(1.1)	(0.2)	(0.0)	(100.0)				
記載なし	76	16	48	150	360	551	648	739	706	544	223	79	13	3	4,156	19,016	23,172	159.51	13.42
(%)	(1.8)	(0.4)	(1.2)	(3.6)	(8.7)	(13.3)	(15.6)	(17.8)	(17.0)	(13.1)	(5.4)	(1.9)	(0.3)	(0.1)	(100.0)				
総計	237	568	2,305	8,283	18,819	30,014	34,857	42,258	37,576	23,827	8,549	2,254	351	57	209,955	19,016	228,971	159.94	9.59
(%)	(0.1)	(0.3)	(1.1)	(3.9)	(9.0)	(14.3)	(16.6)	(20.1)	(17.9)	(11.3)	(4.1)	(1.1)	(0.2)	(0.0)	(100.0)				
平均	23.95	21.33	21.08	20.84	20.79	20.91	21.07	21.50	21.83	22.21	22.67	23.34	23.63	23.60	21.46		21.46		
標準偏差	5.03	3.93	3.70	3.68	3.66	3.70	3.53	3.45	3.55	3.71	3.94	4.18	4.21	6.06	3.66		3.66		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査