

表 2343 身長(cm) 透析前BUN(mg/dL)別（施設血液透析患者 透析回数週3回 透析歴2年以上）

透析前BUN	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
20未満	1	2	9	24	47	78	59	77	71	44	12	5			429	50	479	158.29	10.11
(%)	(0.2)	(0.5)	(2.1)	(5.6)	(11.0)	(18.2)	(13.8)	(17.9)	(16.6)	(10.3)	(2.8)	(1.2)			(100.0)				
20～	7	15	43	139	256	376	407	425	332	171	59	10			2,240	240	2,480	157.35	9.86
(%)	(0.3)	(0.7)	(1.9)	(6.2)	(11.4)	(16.8)	(18.2)	(19.0)	(14.8)	(7.6)	(2.6)	(0.4)			(100.0)				
30～	15	50	174	518	1,065	1,492	1,604	1,907	1,554	872	261	77	2	3	9,594	764	10,358	158.27	9.77
(%)	(0.2)	(0.5)	(1.8)	(5.4)	(11.1)	(15.6)	(16.7)	(19.9)	(16.2)	(9.1)	(2.7)	(0.8)	(0.0)	(0.0)	(100.0)				
40～	39	83	311	1,150	2,505	3,765	4,352	5,116	4,255	2,503	912	220	37	3	25,251	1,560	26,811	159.19	9.59
(%)	(0.2)	(0.3)	(1.2)	(4.6)	(9.9)	(14.9)	(17.2)	(20.3)	(16.9)	(9.9)	(3.6)	(0.9)	(0.1)	(0.0)	(100.0)				
50～	40	97	447	1,565	3,623	5,907	6,784	8,252	7,139	4,459	1,564	397	63	5	40,342	2,317	42,659	159.84	9.45
(%)	(0.1)	(0.2)	(1.1)	(3.9)	(9.0)	(14.6)	(16.8)	(20.5)	(17.7)	(11.1)	(3.9)	(1.0)	(0.2)	(0.0)	(100.0)				
60～	40	68	378	1,398	3,481	5,757	6,767	8,069	7,392	4,844	1,727	439	87	12	40,459	2,184	42,643	160.32	9.46
(%)	(0.1)	(0.2)	(0.9)	(3.5)	(8.6)	(14.2)	(16.7)	(19.9)	(18.3)	(12.0)	(4.3)	(1.1)	(0.2)	(0.0)	(100.0)				
70～	25	57	224	897	2,264	3,723	4,325	5,280	5,152	3,346	1,202	327	40	7	26,869	1,453	28,322	160.58	9.48
(%)	(0.1)	(0.2)	(0.8)	(3.3)	(8.4)	(13.9)	(16.1)	(19.7)	(19.2)	(12.5)	(4.5)	(1.2)	(0.1)	(0.0)	(100.0)				
80～	17	25	108	415	1,090	1,728	2,072	2,472	2,292	1,536	560	155	21	1	12,492	720	13,212	160.43	9.53
(%)	(0.1)	(0.2)	(0.9)	(3.3)	(8.7)	(13.8)	(16.6)	(19.8)	(18.3)	(12.3)	(4.5)	(1.2)	(0.2)	(0.0)	(100.0)				
90～	6	9	45	154	372	652	744	886	794	562	203	63	11	4	4,505	298	4,803	160.48	9.74
(%)	(0.1)	(0.2)	(1.0)	(3.4)	(8.3)	(14.5)	(16.5)	(19.7)	(17.6)	(12.5)	(4.5)	(1.4)	(0.2)	(0.1)	(100.0)				
100～	2	4	9	35	99	161	210	227	240	173	60	24	5		1,249	99	1,348	161.15	9.79
(%)	(0.2)	(0.3)	(0.7)	(2.8)	(7.9)	(12.9)	(16.8)	(18.2)	(19.2)	(13.9)	(4.8)	(1.9)	(0.4)		(100.0)				
110～		1	2	19	30	53	67	77	52	51	17	7			376	18	394	160.18	9.66
(%)		(0.3)	(0.5)	(5.1)	(8.0)	(14.1)	(17.8)	(20.5)	(13.8)	(13.6)	(4.5)	(1.9)			(100.0)				
120～		1	1	5	12	45	26	31	29	23	10	2		1	186	18	204	160.35	9.74
(%)		(0.5)	(0.5)	(2.7)	(6.5)	(24.2)	(14.0)	(16.7)	(15.6)	(12.4)	(5.4)	(1.1)		(0.5)	(100.0)				
合計	192	412	1,751	6,319	14,844	23,737	27,417	32,819	29,302	18,584	6,587	1,726	266	36	163,992	9,721	173,713	159.92	9.55
(%)	(0.1)	(0.3)	(1.1)	(3.9)	(9.1)	(14.5)	(16.7)	(20.0)	(17.9)	(11.3)	(4.0)	(1.1)	(0.2)	(0.0)	(100.0)				
記載なし	1	7	20	75	197	290	311	366	329	199	67	18	1		1,881	2,615	4,496	159.36	9.48
(%)	(0.1)	(0.4)	(1.1)	(4.0)	(10.5)	(15.4)	(16.5)	(19.5)	(17.5)	(10.6)	(3.6)	(1.0)	(0.1)		(100.0)				
総計	193	419	1,771	6,394	15,041	24,027	27,728	33,185	29,631	18,783	6,654	1,744	267	36	165,873	12,336	178,209	159.92	9.55
(%)	(0.1)	(0.3)	(1.1)	(3.9)	(9.1)	(14.5)	(16.7)	(20.0)	(17.9)	(11.3)	(4.0)	(1.1)	(0.2)	(0.0)	(100.0)				
平均	58.71	56.72	57.71	58.73	59.84	60.51	60.79	60.85	61.61	62.49	62.80	63.48	64.26	65.69	61.01	59.86	60.94		
標準偏差	17.36	17.34	16.13	16.09	15.77	15.80	15.53	15.40	15.27	15.33	15.23	15.70	14.02	18.84	15.55	17.00	15.64		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査