

表 2355 身長(cm) %クレアチニン産生速度(%)別（施設血液透析患者 男性のみ 透析回数週3回 透析歴2年以上）

%クレアチニン 産生速度	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
60未満	19	12	12	58	137	461	1,080	1,759	1,685	1,029	363	87	15	4	6,721	703	7,424	163.45	8.20
(%)	(0.3)	(0.2)	(0.2)	(0.9)	(2.0)	(6.9)	(16.1)	(26.2)	(25.1)	(15.3)	(5.4)	(1.3)	(0.2)	(0.1)	(100.0)				
60～	11	4	5	27	88	328	820	1,438	1,352	894	295	85	14	2	5,363	423	5,786	164.05	7.69
(%)	(0.2)	(0.1)	(0.1)	(0.5)	(1.6)	(6.1)	(15.3)	(26.8)	(25.2)	(16.7)	(5.5)	(1.6)	(0.3)	(0.0)	(100.0)				
70～	7	3	10	39	115	473	1,250	2,196	2,150	1,393	497	147	30	3	8,313	596	8,909	164.44	7.38
(%)	(0.1)	(0.0)	(0.1)	(0.5)	(1.4)	(5.7)	(15.0)	(26.4)	(25.9)	(16.8)	(6.0)	(1.8)	(0.4)	(0.0)	(100.0)				
80～	12	8	18	37	140	653	1,682	3,142	3,086	2,033	800	189	37	7	11,844	715	12,559	164.68	7.54
(%)	(0.1)	(0.1)	(0.2)	(0.3)	(1.2)	(5.5)	(14.2)	(26.5)	(26.1)	(17.2)	(6.8)	(1.6)	(0.3)	(0.1)	(100.0)				
90～	6	4	4	40	193	779	2,083	3,912	4,077	2,722	1,017	274	38	6	15,155	834	15,989	164.95	7.28
(%)	(0.0)	(0.0)	(0.0)	(0.3)	(1.3)	(5.1)	(13.7)	(25.8)	(26.9)	(18.0)	(6.7)	(1.8)	(0.3)	(0.0)	(100.0)				
100～	4	2	8	42	157	770	2,121	4,212	4,375	2,900	1,058	304	39	6	15,998	820	16,818	165.13	7.05
(%)	(0.0)	(0.0)	(0.1)	(0.3)	(1.0)	(4.8)	(13.3)	(26.3)	(27.3)	(18.1)	(6.6)	(1.9)	(0.2)	(0.0)	(100.0)				
110～	3		12	21	126	628	1,904	3,757	3,945	2,749	965	254	36	1	14,401	662	15,063	165.27	6.91
(%)	(0.0)		(0.1)	(0.1)	(0.9)	(4.4)	(13.2)	(26.1)	(27.4)	(19.1)	(6.7)	(1.8)	(0.2)	(0.0)	(100.0)				
120～	2		3	14	84	481	1,351	2,768	2,957	1,919	611	168	26	1	10,385	434	10,819	165.15	6.79
(%)	(0.0)		(0.0)	(0.1)	(0.8)	(4.6)	(13.0)	(26.7)	(28.5)	(18.5)	(5.9)	(1.6)	(0.3)	(0.0)	(100.0)				
130～				4	57	257	791	1,590	1,660	1,029	379	70	11	1	5,849	250	6,099	165.06	6.64
(%)				(0.1)	(1.0)	(4.4)	(13.5)	(27.2)	(28.4)	(17.6)	(6.5)	(1.2)	(0.2)	(0.0)	(100.0)				
140～			1	6	50	211	649	1,079	1,101	642	205	46	4	1	3,995	156	4,151	164.36	6.69
(%)			(0.0)	(0.2)	(1.3)	(5.3)	(16.2)	(27.0)	(27.6)	(16.1)	(5.1)	(1.2)	(0.1)	(0.0)	(100.0)				
合計	64	33	73	288	1,147	5,041	13,731	25,853	26,388	17,310	6,190	1,624	250	32	98,024	5,593	103,617	164.80	7.23
(%)	(0.1)	(0.0)	(0.1)	(0.3)	(1.2)	(5.1)	(14.0)	(26.4)	(26.9)	(17.7)	(6.3)	(1.7)	(0.3)	(0.0)	(100.0)				
記載なし	4	6	5	22	80	356	948	1,848	1,891	1,217	407	113	14	3	6,914	1,984	8,898	164.70	7.11
(%)	(0.1)	(0.1)	(0.1)	(0.3)	(1.2)	(5.1)	(13.7)	(26.7)	(27.4)	(17.6)	(5.9)	(1.6)	(0.2)	(0.0)	(100.0)				
総計	68	39	78	310	1,227	5,397	14,679	27,701	28,279	18,527	6,597	1,737	264	35	104,938	7,577	112,515	164.79	7.22
(%)	(0.1)	(0.0)	(0.1)	(0.3)	(1.2)	(5.1)	(14.0)	(26.4)	(26.9)	(17.7)	(6.3)	(1.7)	(0.3)	(0.0)	(100.0)				
平均	73.47	70.04	84.49	83.77	93.93	96.40	98.33	99.50	100.32	100.26	99.71	99.05	96.80	88.76	99.38	92.34	99.00		
標準偏差	25.17	22.13	27.14	27.73	27.81	26.75	26.47	25.58	25.27	24.63	24.06	23.36	22.78	23.97	25.48	27.55	25.64		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査