

表 2360 身長(cm) 透析前リン濃度(mg/dL)別 (施設血液透析患者 透析回数週3回)

透析前リン濃度	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
3未満	24	50	151	487	912	1,355	1,384	1,550	1,202	638	212	40	6	2	8,013	948	8,961	157.57	9.92
(%)	(0.3)	(0.6)	(1.9)	(6.1)	(11.4)	(16.9)	(17.3)	(19.3)	(15.0)	(8.0)	(2.6)	(0.5)	(0.1)	(0.0)	(100.0)				
3.0～	17	47	145	539	1,072	1,526	1,723	2,091	1,638	889	253	72	5	3	10,020	960	10,980	158.39	9.58
(%)	(0.2)	(0.5)	(1.4)	(5.4)	(10.7)	(15.2)	(17.2)	(20.9)	(16.3)	(8.9)	(2.5)	(0.7)	(0.0)	(0.0)	(100.0)				
3.5～	21	66	226	871	1,800	2,662	3,017	3,603	2,959	1,691	514	137	16	2	17,585	1,433	19,018	158.85	9.52
(%)	(0.1)	(0.4)	(1.3)	(5.0)	(10.2)	(15.1)	(17.2)	(20.5)	(16.8)	(9.6)	(2.9)	(0.8)	(0.1)	(0.0)	(100.0)				
4.0～	34	68	355	1,178	2,586	3,903	4,475	5,306	4,594	2,637	856	212	35	1	26,240	1,952	28,192	159.22	9.47
(%)	(0.1)	(0.3)	(1.4)	(4.5)	(9.9)	(14.9)	(17.1)	(20.2)	(17.5)	(10.0)	(3.3)	(0.8)	(0.1)	(0.0)	(100.0)				
4.5～	30	86	355	1,255	2,886	4,664	5,404	6,257	5,420	3,426	1,161	265	38	9	31,256	2,212	33,468	159.61	9.48
(%)	(0.1)	(0.3)	(1.1)	(4.0)	(9.2)	(14.9)	(17.3)	(20.0)	(17.3)	(11.0)	(3.7)	(0.8)	(0.1)	(0.0)	(100.0)				
5.0～	30	85	334	1,242	2,789	4,625	5,171	6,458	5,745	3,546	1,239	325	53	8	31,650	2,159	33,809	159.95	9.50
(%)	(0.1)	(0.3)	(1.1)	(3.9)	(8.8)	(14.6)	(16.3)	(20.4)	(18.2)	(11.2)	(3.9)	(1.0)	(0.2)	(0.0)	(100.0)				
5.5～	24	59	265	926	2,332	3,833	4,525	5,474	4,986	3,188	1,244	315	49	13	27,233	1,779	29,012	160.37	9.49
(%)	(0.1)	(0.2)	(1.0)	(3.4)	(8.6)	(14.1)	(16.6)	(20.1)	(18.3)	(11.7)	(4.6)	(1.2)	(0.2)	(0.0)	(100.0)				
6.0～	5	9	36	158	392	652	749	924	866	530	227	63	10	3	4,624	304	4,928	160.54	9.56
(%)	(0.1)	(0.2)	(0.8)	(3.4)	(8.5)	(14.1)	(16.2)	(20.0)	(18.7)	(11.5)	(4.9)	(1.4)	(0.2)	(0.1)	(100.0)				
6.1～	12	28	117	503	1,241	1,966	2,505	3,126	2,928	1,999	743	205	36	3	15,412	1,093	16,505	160.92	9.48
(%)	(0.1)	(0.2)	(0.8)	(3.3)	(8.1)	(12.8)	(16.3)	(20.3)	(19.0)	(13.0)	(4.8)	(1.3)	(0.2)	(0.0)	(100.0)				
6.5～	11	25	100	367	1,035	1,766	2,157	2,654	2,627	1,804	690	192	29	3	13,460	917	14,377	161.17	9.45
(%)	(0.1)	(0.2)	(0.7)	(2.7)	(7.7)	(13.1)	(16.0)	(19.7)	(19.5)	(13.4)	(5.1)	(1.4)	(0.2)	(0.0)	(100.0)				
7.0～	13	16	75	263	599	1,011	1,307	1,652	1,549	1,198	434	125	23	1	8,266	594	8,860	161.33	9.63
(%)	(0.2)	(0.2)	(0.9)	(3.2)	(7.2)	(12.2)	(15.8)	(20.0)	(18.7)	(14.5)	(5.3)	(1.5)	(0.3)	(0.0)	(100.0)				
7.5～	5	7	49	140	361	600	791	984	983	704	308	88	12	2	5,034	325	5,359	161.64	9.71
(%)	(0.1)	(0.1)	(1.0)	(2.8)	(7.2)	(11.9)	(15.7)	(19.5)	(19.5)	(14.0)	(6.1)	(1.7)	(0.2)	(0.0)	(100.0)				
8.0～	4	5	24	96	193	358	431	578	565	461	200	58	17	2	2,992	200	3,192	162.08	9.89
(%)	(0.1)	(0.2)	(0.8)	(3.2)	(6.5)	(12.0)	(14.4)	(19.3)	(18.9)	(15.4)	(6.7)	(1.9)	(0.6)	(0.1)	(100.0)				
8.5～		2	9	49	126	217	269	346	375	288	125	47	8	4	1,865	158	2,023	162.48	9.65
(%)		(0.1)	(0.5)	(2.6)	(6.8)	(11.6)	(14.4)	(18.6)	(20.1)	(15.4)	(6.7)	(2.5)	(0.4)	(0.2)	(100.0)				
9.0～	5	4	22	75	183	341	411	567	578	463	211	76	12	1	2,949	207	3,156	162.50	9.82
(%)	(0.2)	(0.1)	(0.7)	(2.5)	(6.2)	(11.6)	(13.9)	(19.2)	(19.6)	(15.7)	(7.2)	(2.6)	(0.4)	(0.0)	(100.0)				
合計	235	557	2,263	8,149	18,507	29,479	34,319	41,570	37,015	23,462	8,417	2,220	349	57	206,599	15,241	221,840	159.95	9.59
(%)	(0.1)	(0.3)	(1.1)	(3.9)	(9.0)	(14.3)	(16.6)	(20.1)	(17.9)	(11.4)	(4.1)	(1.1)	(0.2)	(0.0)	(100.0)				
記載なし	2	11	42	134	312	535	538	688	561	365	132	34	2		3,356	3,775	7,131	159.49	9.52
(%)	(0.1)	(0.3)	(1.3)	(4.0)	(9.3)	(15.9)	(16.0)	(20.5)	(16.7)	(10.9)	(3.9)	(1.0)	(0.1)		(100.0)				
総計	237	568	2,305	8,283	18,819	30,014	34,857	42,258	37,576	23,827	8,549	2,254	351	57	209,955	19,016	228,971	159.94	9.59
(%)	(0.1)	(0.3)	(1.1)	(3.9)	(9.0)	(14.3)	(16.6)	(20.1)	(17.9)	(11.3)	(4.1)	(1.1)	(0.2)	(0.0)	(100.0)				
平均	4.90	4.74	4.90	4.93	5.03	5.10	5.15	5.20	5.29	5.42	5.56	5.70	5.85	5.68	5.21	5.06	5.20		
標準偏差	1.63	1.39	1.42	1.39	1.38	1.39	1.39	1.40	1.42	1.45	1.51	1.58	1.58	1.65	1.42	1.50	1.43		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査