

表 2362 身長(cm) PTH値(pg/mL)別（施設血液透析患者 透析回数週3回）PTH測定法:whole-PTH

PTH値	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
20未満	2	7	12	64	148	197	257	299	232	137	41	15	1		1,412	87	1,499	159.05	9.68
(%)	(0.1)	(0.5)	(0.8)	(4.5)	(10.5)	(14.0)	(18.2)	(21.2)	(16.4)	(9.7)	(2.9)	(1.1)	(0.1)		(100.0)				
20～	4	4	17	73	176	282	297	344	277	190	66	18	2		1,750	98	1,848	159.22	9.56
(%)	(0.2)	(0.2)	(1.0)	(4.2)	(10.1)	(16.1)	(17.0)	(19.7)	(15.8)	(10.9)	(3.8)	(1.0)	(0.1)		(100.0)				
35～	3	3	19	65	147	205	259	311	273	154	57	11	1		1,508	105	1,613	159.52	9.76
(%)	(0.2)	(0.2)	(1.3)	(4.3)	(9.7)	(13.6)	(17.2)	(20.6)	(18.1)	(10.2)	(3.8)	(0.7)	(0.1)		(100.0)				
50～	2	6	28	92	191	291	394	483	421	255	121	34	5	1	2,324	135	2,459	160.42	9.89
(%)	(0.1)	(0.3)	(1.2)	(4.0)	(8.2)	(12.5)	(17.0)	(20.8)	(18.1)	(11.0)	(5.2)	(1.5)	(0.2)	(0.0)	(100.0)				
75～	3	6	17	54	176	248	351	429	402	240	77	20	5	1	2,029	119	2,148	160.60	9.42
(%)	(0.1)	(0.3)	(0.8)	(2.7)	(8.7)	(12.2)	(17.3)	(21.1)	(19.8)	(11.8)	(3.8)	(1.0)	(0.2)	(0.0)	(100.0)				
100～		3	18	56	132	221	255	323	383	208	79	18	1		1,697	90	1,787	160.93	9.30
(%)		(0.2)	(1.1)	(3.3)	(7.8)	(13.0)	(15.0)	(19.0)	(22.6)	(12.3)	(4.7)	(1.1)	(0.1)		(100.0)				
125～	1	3	8	33	86	142	154	213	233	155	57	18	1		1,104	83	1,187	161.37	9.47
(%)	(0.1)	(0.3)	(0.7)	(3.0)	(7.8)	(12.9)	(13.9)	(19.3)	(21.1)	(14.0)	(5.2)	(1.6)	(0.1)		(100.0)				
150～			1	2	3	4	4	9	7	4	1	1			36	1	37	160.11	10.12
(%)			(2.8)	(5.6)	(8.3)	(11.1)	(11.1)	(25.0)	(19.4)	(11.1)	(2.8)	(2.8)			(100.0)				
151～	1	3	7	33	75	121	125	168	171	121	34	10	1		870	64	934	160.57	9.82
(%)	(0.1)	(0.3)	(0.8)	(3.8)	(8.6)	(13.9)	(14.4)	(19.3)	(19.7)	(13.9)	(3.9)	(1.1)	(0.1)		(100.0)				
180～			4	35	49	76	109	106	131	75	26	11	1		623	43	666	160.75	9.58
(%)			(0.6)	(5.6)	(7.9)	(12.2)	(17.5)	(17.0)	(21.0)	(12.0)	(4.2)	(1.8)	(0.2)		(100.0)				
210～			6	17	34	52	74	82	85	53	31	2	2		438	34	472	160.95	9.49
(%)			(1.4)	(3.9)	(7.8)	(11.9)	(16.9)	(18.7)	(19.4)	(12.1)	(7.1)	(0.5)	(0.5)		(100.0)				
240～		3	5	16	45	72	70	83	83	54	23	8	3		465	61	526	160.29	10.07
(%)		(0.6)	(1.1)	(3.4)	(9.7)	(15.5)	(15.1)	(17.8)	(17.8)	(11.6)	(4.9)	(1.7)	(0.6)		(100.0)				
300～			3	9	15	19	28	43	27	25	15	4			188	17	205	161.36	10.05
(%)			(1.6)	(4.8)	(8.0)	(10.1)	(14.9)	(22.9)	(14.4)	(13.3)	(8.0)	(2.1)			(100.0)				
350～			3	7	12	14	14	20	18	20	5	1			114	21	135	159.86	10.50
(%)			(2.6)	(6.1)	(10.5)	(12.3)	(12.3)	(17.5)	(15.8)	(17.5)	(4.4)	(0.9)			(100.0)				
400～	1	1	2	5	16	19	21	18	28	15	10	4			140	12	152	160.05	11.21
(%)	(0.7)	(0.7)	(1.4)	(3.6)	(11.4)	(13.6)	(15.0)	(12.9)	(20.0)	(10.7)	(7.1)	(2.9)			(100.0)				
500～		1	3	9	13	17	31	23	28	21	9	1			156	10	166	160.21	10.28
(%)		(0.6)	(1.9)	(5.8)	(8.3)	(10.9)	(19.9)	(14.7)	(17.9)	(13.5)	(5.8)	(0.6)			(100.0)				
合計	17	40	153	570	1,318	1,980	2,443	2,954	2,799	1,727	652	176	23	2	14,854	980	15,834	160.25	9.68
(%)	(0.1)	(0.3)	(1.0)	(3.8)	(8.9)	(13.3)	(16.4)	(19.9)	(18.8)	(11.6)	(4.4)	(1.2)	(0.2)	(0.0)	(100.0)				
記載なし	1	2	4	21	29	56	57	62	56	37	13	5			343	95	438	159.01	10.07
(%)	(0.3)	(0.6)	(1.2)	(6.1)	(8.5)	(16.3)	(16.6)	(18.1)	(16.3)	(10.8)	(3.8)	(1.5)			(100.0)				
総計	18	42	157	591	1,347	2,036	2,500	3,016	2,855	1,764	665	181	23	2	15,197	1,075	16,272	160.22	9.69
(%)	(0.1)	(0.3)	(1.0)	(3.9)	(8.9)	(13.4)	(16.5)	(19.8)	(18.8)	(11.6)	(4.4)	(1.2)	(0.2)	(0.0)	(100.0)				
平均	83.59	107.55	128.47	108.40	100.79	103.22	101.33	99.51	107.18	110.78	117.25	114.73	114.43	70.00	104.80	119.37	105.70		
標準偏差	103.21	122.77	224.24	118.75	100.27	130.45	101.78	97.47	104.09	104.40	114.56	103.38	81.31	19.80	109.11	110.40	109.24		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査