

表 2363 身長(cm) 透析前ヘモグロビン濃度(g/dL)別（施設血液透析患者 透析回数週3回）

透析前ヘモグロビン濃度	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
5未満		2	1	3	27	41	37	54	42	22	11	1			241	20	261	159.45	9.09
(%)		(0.8)	(0.4)	(1.2)	(11.2)	(17.0)	(15.4)	(22.4)	(17.4)	(9.1)	(4.6)	(0.4)			(100.0)				
5～	1		2	20	29	62	59	59	50	30	14		1		327	35	362	158.75	9.39
(%)	(0.3)		(0.6)	(6.1)	(8.9)	(19.0)	(18.0)	(18.0)	(15.3)	(9.2)	(4.3)		(0.3)		(100.0)				
6～	2	1	14	43	85	125	151	156	152	72	31	9	2		843	99	942	158.79	10.21
(%)	(0.2)	(0.1)	(1.7)	(5.1)	(10.1)	(14.8)	(17.9)	(18.5)	(18.0)	(8.5)	(3.7)	(1.1)	(0.2)		(100.0)				
7～	6	17	47	167	338	537	565	600	502	316	96	26	9	1	3,227	351	3,578	158.48	9.88
(%)	(0.2)	(0.5)	(1.5)	(5.2)	(10.5)	(16.6)	(17.5)	(18.6)	(15.6)	(9.8)	(3.0)	(0.8)	(0.3)	(0.0)	(100.0)				
8～	20	59	201	571	1,138	1,687	1,812	2,087	1,708	1,020	343	86	14	1	10,747	979	11,726	158.45	9.90
(%)	(0.2)	(0.5)	(1.9)	(5.3)	(10.6)	(15.7)	(16.9)	(19.4)	(15.9)	(9.5)	(3.2)	(0.8)	(0.1)	(0.0)	(100.0)				
9～	43	95	436	1,590	3,571	5,495	5,979	6,708	5,695	3,408	1,123	277	40	13	34,473	2,565	37,038	158.94	9.54
(%)	(0.1)	(0.3)	(1.3)	(4.6)	(10.4)	(15.9)	(17.3)	(19.5)	(16.5)	(9.9)	(3.3)	(0.8)	(0.1)	(0.0)	(100.0)				
10～	78	174	752	2,790	6,449	10,374	11,824	14,250	12,435	7,796	2,788	723	110	15	70,558	4,659	75,217	159.81	9.53
(%)	(0.1)	(0.2)	(1.1)	(4.0)	(9.1)	(14.7)	(16.8)	(20.2)	(17.6)	(11.0)	(4.0)	(1.0)	(0.2)	(0.0)	(100.0)				
11～	54	144	539	2,013	4,778	7,755	9,364	11,838	10,734	6,977	2,613	678	111	21	57,619	3,860	61,479	160.52	9.52
(%)	(0.1)	(0.2)	(0.9)	(3.5)	(8.3)	(13.5)	(16.3)	(20.5)	(18.6)	(12.1)	(4.5)	(1.2)	(0.2)	(0.0)	(100.0)				
12～	19	38	211	683	1,571	2,487	3,356	4,264	4,099	2,680	957	295	42	3	20,705	1,665	22,370	161.00	9.45
(%)	(0.1)	(0.2)	(1.0)	(3.3)	(7.6)	(12.0)	(16.2)	(20.6)	(19.8)	(12.9)	(4.6)	(1.4)	(0.2)	(0.0)	(100.0)				
13～	5	15	46	175	350	563	760	1,042	1,052	710	263	82	10		5,073	464	5,537	161.45	9.61
(%)	(0.1)	(0.3)	(0.9)	(3.4)	(6.9)	(11.1)	(15.0)	(20.5)	(20.7)	(14.0)	(5.2)	(1.6)	(0.2)		(100.0)				
14～	5	6	11	42	95	179	245	356	369	297	129	37	7	3	1,781	152	1,933	162.78	10.02
(%)	(0.3)	(0.3)	(0.6)	(2.4)	(5.3)	(10.1)	(13.8)	(20.0)	(20.7)	(16.7)	(7.2)	(2.1)	(0.4)	(0.2)	(100.0)				
合計	233	551	2,260	8,097	18,431	29,305	34,152	41,414	36,838	23,328	8,368	2,214	346	57	205,594	14,849	220,443	159.95	9.59
(%)	(0.1)	(0.3)	(1.1)	(3.9)	(9.0)	(14.3)	(16.6)	(20.1)	(17.9)	(11.3)	(4.1)	(1.1)	(0.2)	(0.0)	(100.0)				
記載なし	4	17	45	186	388	709	705	844	738	499	181	40	5		4,361	4,167	8,528	159.57	9.61
(%)	(0.1)	(0.4)	(1.0)	(4.3)	(8.9)	(16.3)	(16.2)	(19.4)	(16.9)	(11.4)	(4.2)	(0.9)	(0.1)		(100.0)				
総計	237	568	2,305	8,283	18,819	30,014	34,857	42,258	37,576	23,827	8,549	2,254	351	57	209,955	19,016	228,971	159.94	9.59
(%)	(0.1)	(0.3)	(1.1)	(3.9)	(9.0)	(14.3)	(16.6)	(20.1)	(17.9)	(11.3)	(4.1)	(1.1)	(0.2)	(0.0)	(100.0)				
平均	10.49	10.47	10.49	10.52	10.55	10.57	10.65	10.72	10.77	10.83	10.87	10.96	10.88	10.81	10.69	10.65	10.69		
標準偏差	1.39	1.41	1.30	1.27	1.25	1.25	1.26	1.26	1.27	1.28	1.30	1.28	1.36	1.30	1.27	1.40	1.28		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査