

表 2364 身長(cm) 総コレステロール濃度(mg/dL)別（施設血液透析患者 透析回数週3回）

総コレステロール濃度	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
100未満	10	10	46	132	327	535	943	1,397	1,284	801	305	86	10	5	5,891	563	6,454	162.12	9.06
(%)	(0.2)	(0.2)	(0.8)	(2.2)	(5.6)	(9.1)	(16.0)	(23.7)	(21.8)	(13.6)	(5.2)	(1.5)	(0.2)	(0.1)	(100.0)				
100～	20	37	122	501	1,059	1,827	2,666	3,984	3,796	2,264	804	208	29	7	17,324	1,296	18,620	161.63	9.10
(%)	(0.1)	(0.2)	(0.7)	(2.9)	(6.1)	(10.5)	(15.4)	(23.0)	(21.9)	(13.1)	(4.6)	(1.2)	(0.2)	(0.0)	(100.0)				
120～	29	82	307	1,096	2,289	3,833	5,215	6,998	6,416	3,910	1,450	378	59	9	32,071	2,244	34,315	160.91	9.35
(%)	(0.1)	(0.3)	(1.0)	(3.4)	(7.1)	(12.0)	(16.3)	(21.8)	(20.0)	(12.2)	(4.5)	(1.2)	(0.2)	(0.0)	(100.0)				
140～	40	95	427	1,418	3,125	4,913	5,774	7,472	6,758	4,279	1,491	387	67	13	36,259	2,567	38,826	160.20	9.61
(%)	(0.1)	(0.3)	(1.2)	(3.9)	(8.6)	(13.5)	(15.9)	(20.6)	(18.6)	(11.8)	(4.1)	(1.1)	(0.2)	(0.0)	(100.0)				
160～	42	78	363	1,323	3,039	4,573	4,971	5,660	4,959	3,211	1,139	302	45	4	29,709	1,874	31,583	159.34	9.73
(%)	(0.1)	(0.3)	(1.2)	(4.5)	(10.2)	(15.4)	(16.7)	(19.1)	(16.7)	(10.8)	(3.8)	(1.0)	(0.2)	(0.0)	(100.0)				
180～	21	58	251	896	2,165	3,161	3,249	3,242	2,712	1,777	627	174	29	3	18,365	1,254	19,619	158.48	9.71
(%)	(0.1)	(0.3)	(1.4)	(4.9)	(11.8)	(17.2)	(17.7)	(17.7)	(14.8)	(9.7)	(3.4)	(0.9)	(0.2)	(0.0)	(100.0)				
200～	10	38	134	498	1,155	1,900	1,787	1,581	1,267	827	297	74	17	1	9,586	623	10,209	157.74	9.62
(%)	(0.1)	(0.4)	(1.4)	(5.2)	(12.0)	(19.8)	(18.6)	(16.5)	(13.2)	(8.6)	(3.1)	(0.8)	(0.2)	(0.0)	(100.0)				
220～	4	12	49	221	553	854	779	691	534	332	104	31	3		4,167	308	4,475	157.37	9.32
(%)	(0.1)	(0.3)	(1.2)	(5.3)	(13.3)	(20.5)	(18.7)	(16.6)	(12.8)	(8.0)	(2.5)	(0.7)	(0.1)		(100.0)				
240～	2	8	21	83	239	357	328	236	154	114	49	10	3		1,604	129	1,733	156.59	9.32
(%)	(0.1)	(0.5)	(1.3)	(5.2)	(14.9)	(22.3)	(20.4)	(14.7)	(9.6)	(7.1)	(3.1)	(0.6)	(0.2)		(100.0)				
260～		1	21	64	136	230	197	141	110	78	29	10	1		1,018	93	1,111	156.78	9.54
(%)		(0.1)	(2.1)	(6.3)	(13.4)	(22.6)	(19.4)	(13.9)	(10.8)	(7.7)	(2.8)	(1.0)	(0.1)		(100.0)				
合計	178	419	1,741	6,232	14,087	22,183	25,909	31,402	27,990	17,593	6,295	1,660	263	42	155,994	10,951	166,945	159.92	9.59
(%)	(0.1)	(0.3)	(1.1)	(4.0)	(9.0)	(14.2)	(16.6)	(20.1)	(17.9)	(11.3)	(4.0)	(1.1)	(0.2)	(0.0)	(100.0)				
記載なし	59	149	564	2,051	4,732	7,831	8,948	10,856	9,586	6,234	2,254	594	88	15	53,961	8,065	62,026	160.00	9.57
(%)	(0.1)	(0.3)	(1.0)	(3.8)	(8.8)	(14.5)	(16.6)	(20.1)	(17.8)	(11.6)	(4.2)	(1.1)	(0.2)	(0.0)	(100.0)				
総計	237	568	2,305	8,283	18,819	30,014	34,857	42,258	37,576	23,827	8,549	2,254	351	57	209,955	19,016	228,971	159.94	9.59
(%)	(0.1)	(0.3)	(1.1)	(3.9)	(9.0)	(14.3)	(16.6)	(20.1)	(17.9)	(11.3)	(4.1)	(1.1)	(0.2)	(0.0)	(100.0)				
平均	153.91	160.15	161.00	161.78	163.29	162.25	156.80	150.99	149.35	150.26	149.92	150.10	152.04	138.83	154.81	153.42	154.71		
標準偏差	33.96	35.22	35.40	35.33	35.97	36.43	35.89	34.07	33.23	33.42	33.93	34.54	33.18	28.73	35.16	36.98	35.28		

数値下のかっこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2014年調査