

表 2367 身長(cm) 透析前収縮期血圧(mmHg)別（施設血液透析患者 透析回数週3回）

透析前収縮期血圧	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
60未満	1		4	2	1	9	9	3	1	2	1				33	2	35	152.85	11.97
(%)	(3.0)		(12.1)	(6.1)	(3.0)	(27.3)	(27.3)	(9.1)	(3.0)	(6.1)	(3.0)				(100.0)				
60～		2	3	21	28	47	44	40	37	29	2	1			254	17	271	157.38	9.74
(%)		(0.8)	(1.2)	(8.3)	(11.0)	(18.5)	(17.3)	(15.7)	(14.6)	(11.4)	(0.8)	(0.4)			(100.0)				
80～	7	12	46	124	248	381	399	401	339	150	65	20	2	1	2,195	182	2,377	157.65	9.83
(%)	(0.3)	(0.5)	(2.1)	(5.6)	(11.3)	(17.4)	(18.2)	(18.3)	(15.4)	(6.8)	(3.0)	(0.9)	(0.1)	(0.0)	(100.0)				
100～	17	48	191	639	1,297	1,941	2,127	2,361	1,939	1,171	371	79	16	1	12,198	957	13,155	158.54	9.65
(%)	(0.1)	(0.4)	(1.6)	(5.2)	(10.6)	(15.9)	(17.4)	(19.4)	(15.9)	(9.6)	(3.0)	(0.6)	(0.1)	(0.0)	(100.0)				
120～	43	121	467	1,648	3,645	5,682	6,733	7,783	6,668	4,107	1,423	365	57	7	38,749	2,827	41,576	159.49	9.60
(%)	(0.1)	(0.3)	(1.2)	(4.3)	(9.4)	(14.7)	(17.4)	(20.1)	(17.2)	(10.6)	(3.7)	(0.9)	(0.1)	(0.0)	(100.0)				
140～	71	153	610	2,217	5,237	8,461	9,957	12,386	10,911	6,867	2,399	653	89	10	60,021	4,182	64,203	160.09	9.48
(%)	(0.1)	(0.3)	(1.0)	(3.7)	(8.7)	(14.1)	(16.6)	(20.6)	(18.2)	(11.4)	(4.0)	(1.1)	(0.1)	(0.0)	(100.0)				
160～	43	87	447	1,633	3,701	5,962	7,089	8,861	8,206	5,257	1,948	507	83	12	43,836	3,007	46,843	160.40	9.52
(%)	(0.1)	(0.2)	(1.0)	(3.7)	(8.4)	(13.6)	(16.2)	(20.2)	(18.7)	(12.0)	(4.4)	(1.2)	(0.2)	(0.0)	(100.0)				
180～	21	43	187	663	1,486	2,336	2,571	3,269	3,107	2,015	814	226	32	7	16,777	1,225	18,002	160.37	9.83
(%)	(0.1)	(0.3)	(1.1)	(4.0)	(8.9)	(13.9)	(15.3)	(19.5)	(18.5)	(12.0)	(4.9)	(1.3)	(0.2)	(0.0)	(100.0)				
200～	5	8	36	155	412	580	636	796	752	529	234	61	13	3	4,220	295	4,515	160.56	9.98
(%)	(0.1)	(0.2)	(0.9)	(3.7)	(9.8)	(13.7)	(15.1)	(18.9)	(17.8)	(12.5)	(5.5)	(1.4)	(0.3)	(0.1)	(100.0)				
220～	1	6	9	47	80	111	117	157	134	93	64	13	8		840	54	894	160.22	11.19
(%)	(0.1)	(0.7)	(1.1)	(5.6)	(9.5)	(13.2)	(13.9)	(18.7)	(16.0)	(11.1)	(7.6)	(1.5)	(1.0)		(100.0)				
合計	209	480	2,000	7,149	16,135	25,510	29,682	36,057	32,094	20,220	7,321	1,925	300	41	179,123	12,748	191,871	159.93	9.60
(%)	(0.1)	(0.3)	(1.1)	(4.0)	(9.0)	(14.2)	(16.6)	(20.1)	(17.9)	(11.3)	(4.1)	(1.1)	(0.2)	(0.0)	(100.0)				
記載なし	28	88	305	1,134	2,684	4,504	5,175	6,201	5,482	3,607	1,228	329	51	16	30,832	6,268	37,100	160.00	9.50
(%)	(0.1)	(0.3)	(1.0)	(3.7)	(8.7)	(14.6)	(16.8)	(20.1)	(17.8)	(11.7)	(4.0)	(1.1)	(0.2)	(0.1)	(100.0)				
総計	237	568	2,305	8,283	18,819	30,014	34,857	42,258	37,576	23,827	8,549	2,254	351	57	209,955	19,016	228,971	159.94	9.59
(%)	(0.1)	(0.3)	(1.1)	(3.9)	(9.0)	(14.3)	(16.6)	(20.1)	(17.9)	(11.3)	(4.1)	(1.1)	(0.2)	(0.0)	(100.0)				
平均	149.56	147.71	148.77	149.83	150.62	150.70	150.63	151.64	152.58	153.32	155.04	155.66	157.08	159.32	151.68	151.10	151.64		
標準偏差	27.11	26.04	25.91	25.45	24.88	24.46	23.86	23.51	23.41	23.42	24.02	23.63	26.09	26.25	23.99	24.43	24.02		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査