

表 2373 身長(cm) 原疾患別（血液透析濾過患者 透析回数週3回）

原疾患	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	11	25	93	413	1,129	2,175	2,440	2,671	2,534	1,758	617	164	23	3	14,056	1,020	15,076	160.50	9.31
(%)	(0.1)	(0.2)	(0.7)	(2.9)	(8.0)	(15.5)	(17.4)	(19.0)	(18.0)	(12.5)	(4.4)	(1.2)	(0.2)	(0.0)	(100.0)				
慢性腎盂腎炎	1	2	5	19	46	69	73	63	55	42	22	5	1		403	19	422	159.00	10.17
(%)	(0.2)	(0.5)	(1.2)	(4.7)	(11.4)	(17.1)	(18.1)	(15.6)	(13.6)	(10.4)	(5.5)	(1.2)	(0.2)		(100.0)				
急速進行性糸球体腎炎		1	3	8	23	40	57	51	41	28	7	6	2		267	17	284	159.84	9.58
(%)		(0.4)	(1.1)	(3.0)	(8.6)	(15.0)	(21.3)	(19.1)	(15.4)	(10.5)	(2.6)	(2.2)	(0.7)		(100.0)				
妊娠腎/妊娠中毒症			3	24	68	114	91	31	6						337	22	359	152.53	5.59
(%)			(0.9)	(7.1)	(20.2)	(33.8)	(27.0)	(9.2)	(1.8)						(100.0)				
その他の分類不能の腎炎	1		1	5	10	31	28	29	39	27	13	2			186	9	195	160.93	12.19
(%)	(0.5)		(0.5)	(2.7)	(5.4)	(16.7)	(15.1)	(15.6)	(21.0)	(14.5)	(7.0)	(1.1)			(100.0)				
多発性嚢胞腎			9	27	81	178	235	253	225	207	110	26	4		1,355	109	1,464	162.30	9.38
(%)			(0.7)	(2.0)	(6.0)	(13.1)	(17.3)	(18.7)	(16.6)	(15.3)	(8.1)	(1.9)	(0.3)		(100.0)				
腎硬化症	1	11	49	97	270	418	451	566	436	266	89	21	3		2,678	208	2,886	159.16	9.42
(%)	(0.0)	(0.4)	(1.8)	(3.6)	(10.1)	(15.6)	(16.8)	(21.1)	(16.3)	(9.9)	(3.3)	(0.8)	(0.1)		(100.0)				
悪性高血圧	1	2	3	5	22	36	41	50	58	49	10	10	1		288	24	312	161.66	11.47
(%)	(0.3)	(0.7)	(1.0)	(1.7)	(7.6)	(12.5)	(14.2)	(17.4)	(20.1)	(17.0)	(3.5)	(3.5)	(0.3)		(100.0)				
糖尿病性腎症	13	13	59	353	850	1,376	1,788	2,481	2,413	1,826	675	189	33	6	12,075	1,098	13,173	161.88	9.39
(%)	(0.1)	(0.1)	(0.5)	(2.9)	(7.0)	(11.4)	(14.8)	(20.5)	(20.0)	(15.1)	(5.6)	(1.6)	(0.3)	(0.0)	(100.0)				
SLE腎炎			4	14	43	80	74	51	37	23	5	5			336	27	363	156.97	8.58
(%)			(1.2)	(4.2)	(12.8)	(23.8)	(22.0)	(15.2)	(11.0)	(6.8)	(1.5)	(1.5)			(100.0)				
アミロイド腎				1	9	12	12	9	7	3	2				55	4	59	157.40	8.45
(%)				(1.8)	(16.4)	(21.8)	(21.8)	(16.4)	(12.7)	(5.5)	(3.6)				(100.0)				
痛風腎				3	7	10	20	31	34	27	7	1	2		142	12	154	163.56	8.57
(%)				(2.1)	(4.9)	(7.0)	(14.1)	(21.8)	(23.9)	(19.0)	(4.9)	(0.7)	(1.4)		(100.0)				
先天性代謝異常による腎不全		2	5	2	1	6	4	8	10	3					41	3	44	156.39	12.02
(%)		(4.9)	(12.2)	(4.9)	(2.4)	(14.6)	(9.8)	(19.5)	(24.4)	(7.3)					(100.0)				
腎・尿路結核			1	1	4	5	4	4	1	2					22		22	155.41	8.53
(%)			(4.5)	(4.5)	(18.2)	(22.7)	(18.2)	(18.2)	(4.5)	(9.1)					(100.0)				
腎・尿路結石	1		1	3	4	8	12	12	7	5	7	2			62	7	69	160.48	11.50
(%)	(1.6)		(1.6)	(4.8)	(6.5)	(12.9)	(19.4)	(19.4)	(11.3)	(8.1)	(11.3)	(3.2)			(100.0)				
腎・尿路腫瘍				2	2	9	12	21	23	6	3	2			80	5	85	162.44	7.91
(%)				(2.5)	(2.5)	(11.3)	(15.0)	(26.3)	(28.8)	(7.5)	(3.8)	(2.5)			(100.0)				
閉塞性尿路障害			2	2	7	7	8	10	10	13	4	1	1		65	9	74	161.28	11.15
(%)			(3.1)	(3.1)	(10.8)	(10.8)	(12.3)	(15.4)	(15.4)	(20.0)	(6.2)	(1.5)	(1.5)		(100.0)				
骨髄腫				2		1	5	2	3	4	1				18	1	19	161.33	9.92
(%)				(11.1)		(5.6)	(27.8)	(11.1)	(16.7)	(22.2)	(5.6)				(100.0)				
腎形成不全	1	2	2	9	9	11	19	15	17	8	4	1			98	3	101	157.62	11.21
(%)	(1.0)	(2.0)	(2.0)	(9.2)	(9.2)	(11.2)	(19.4)	(15.3)	(17.3)	(8.2)	(4.1)	(1.0)			(100.0)				
不明	5	9	38	137	308	467	520	561	559	336	139	35	4		3,118	362	3,480	159.53	10.02
(%)	(0.2)	(0.3)	(1.2)	(4.4)	(9.9)	(15.0)	(16.7)	(18.0)	(17.9)	(10.8)	(4.5)	(1.1)	(0.1)		(100.0)				
再導入		1	2	4	24	33	64	95	80	54	27	8	2		394	22	416	162.90	8.99
(%)		(0.3)	(0.5)	(1.0)	(6.1)	(8.4)	(16.2)	(24.1)	(20.3)	(13.7)	(6.9)	(2.0)	(0.5)		(100.0)				
その他	3	2	9	32	78	126	121	163	127	98	44	15	1		819	70	889	160.06	10.01
(%)	(0.4)	(0.2)	(1.1)	(3.9)	(9.5)	(15.4)	(14.8)	(19.9)	(15.5)	(12.0)	(5.4)	(1.8)	(0.1)		(100.0)				
合計	38	70	289	1,163	2,995	5,212	6,079	7,177	6,722	4,785	1,786	493	77	9	36,895	3,051	39,946	160.74	9.54
(%)	(0.1)	(0.2)	(0.8)	(3.2)	(8.1)	(14.1)	(16.5)	(19.5)	(18.2)	(13.0)	(4.8)	(1.3)	(0.2)	(0.0)	(100.0)				
記載なし																			
(%)																			
総計	38	70	289	1,163	2,995	5,212	6,079	7,177	6,722	4,785	1,786	493	77	9	36,895	3,051	39,946	160.74	9.54
(%)	(0.1)	(0.2)	(0.8)	(3.2)	(8.1)	(14.1)	(16.5)	(19.5)	(18.2)	(13.0)	(4.8)	(1.3)	(0.2)	(0.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2014年調査