

表 2375 身長(cm) 透析後体重(kg)別（血液透析濾過患者 透析回数週3回）

透析後体重	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
20未満						3	2	1	1	4					11	1	12	161.73	8.95
(%)						(27.3)	(18.2)	(9.1)	(9.1)	(36.4)					(100.0)				
20～	7	7	10	17	13	8	5	3	2						72	4	76	142.69	10.67
(%)	(9.7)	(9.7)	(13.9)	(23.6)	(18.1)	(11.1)	(6.9)	(4.2)	(2.8)						(100.0)				
30～	15	44	131	414	744	715	294	105	27	4	2	2			2,497	185	2,682	148.57	7.03
(%)	(0.6)	(1.8)	(5.2)	(16.6)	(29.8)	(28.6)	(11.8)	(4.2)	(1.1)	(0.2)	(0.1)	(0.1)			(100.0)				
40～	9	9	107	527	1,458	2,525	2,325	1,555	639	177	26	5			9,362	643	10,005	154.69	7.32
(%)	(0.1)	(0.1)	(1.1)	(5.6)	(15.6)	(27.0)	(24.8)	(16.6)	(6.8)	(1.9)	(0.3)	(0.1)			(100.0)				
50～	5	8	30	155	583	1,406	2,288	3,115	2,575	1,262	278	34	2	1	11,742	748	12,490	161.17	7.65
(%)	(0.0)	(0.1)	(0.3)	(1.3)	(5.0)	(12.0)	(19.5)	(26.5)	(21.9)	(10.7)	(2.4)	(0.3)	(0.0)	(0.0)	(100.0)				
60～	1		4	33	140	418	808	1,689	2,247	1,767	591	127	11		7,836	529	8,365	165.61	7.19
(%)	(0.0)		(0.1)	(0.4)	(1.8)	(5.3)	(10.3)	(21.6)	(28.7)	(22.5)	(7.5)	(1.6)	(0.1)		(100.0)				
70～		1	1	6	26	74	220	490	816	959	463	145	22	3	3,226	197	3,423	168.65	7.20
(%)		(0.0)	(0.0)	(0.2)	(0.8)	(2.3)	(6.8)	(15.2)	(25.3)	(29.7)	(14.4)	(4.5)	(0.7)	(0.1)	(100.0)				
80～			1	10	23	60	116	259	367	367	237	90	17	2	1,182	91	1,273	170.43	7.32
(%)			(0.1)	(0.8)	(1.9)	(5.1)	(9.8)	(21.9)	(31.0)	(20.1)	(7.6)	(1.4)	(0.2)	(0.2)	(100.0)				
90～				2	6	10	27	66	136	113	49	9	2	2	420	35	455	172.55	6.84
(%)				(0.5)	(1.4)	(2.4)	(6.4)	(15.7)	(32.4)	(26.9)	(11.7)	(2.1)	(0.5)	(100.0)					
100～				1		3	4	18	42	35	24	6			133	9	142	174.22	6.69
(%)				(0.8)		(2.3)	(3.0)	(13.5)	(31.6)	(26.3)	(18.0)	(4.5)			(100.0)				
110～				1		2	7	19	15	9	5				58	3	61	174.57	6.64
(%)				(1.7)		(3.4)	(12.1)	(32.8)	(25.9)	(15.5)	(8.6)				(100.0)				
120～				1		1	2	9	12	5	5			1	36	1	37	176.39	7.48
(%)				(2.8)		(2.8)	(5.6)	(25.0)	(33.3)	(13.9)	(13.9)			(2.8)	(100.0)				
合計	37	69	284	1,152	2,977	5,178	6,017	7,108	6,659	4,746	1,772	490	77	9	36,575	2,446	39,021	160.74	9.54
(%)	(0.1)	(0.2)	(0.8)	(3.1)	(8.1)	(14.2)	(16.5)	(19.4)	(18.2)	(13.0)	(4.8)	(1.3)	(0.2)	(0.0)	(100.0)				
記載なし	1	1	5	11	18	34	62	69	63	39	14	3			320	605	925	160.74	9.58
(%)	(0.3)	(0.3)	(1.6)	(3.4)	(5.6)	(10.6)	(19.4)	(21.6)	(19.7)	(12.2)	(4.4)	(0.9)			(100.0)				
総計	38	70	289	1,163	2,995	5,212	6,079	7,177	6,722	4,785	1,786	493	77	9	36,895	3,051	39,946	160.74	9.54
(%)	(0.1)	(0.2)	(0.8)	(3.2)	(8.1)	(14.1)	(16.5)	(19.5)	(18.2)	(13.0)	(4.8)	(1.3)	(0.2)	(0.0)	(100.0)				
平均	39.86	38.31	41.35	43.14	45.64	48.55	52.56	56.87	61.52	66.58	71.89	77.18	86.17	89.22	56.64	56.45	56.63		
標準偏差	9.74	8.46	8.11	7.80	8.43	8.76	9.39	9.58	10.40	12.00	13.68	15.12	18.35	27.65	13.02	13.22	13.03		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査