

表 2388 身長(cm) 透析前クレアチニン濃度(mg/dL)別（血液透析濾過患者 透析回数週3回 透析歴2年以上）

透析前クレアチニン濃度	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
2未満				1		2									3		3	147.00	6.08
(%)				(33.3)		(66.7)									(100.0)				
2～	2		3	15	18	27	19	19	23	4					130	17	147	154.52	10.02
(%)	(1.5)		(2.3)	(11.5)	(13.8)	(20.8)	(14.6)	(14.6)	(17.7)	(3.1)					(100.0)				
4～	3	6	17	74	148	170	144	119	75	45	13	6			820	98	918	154.85	9.54
(%)	(0.4)	(0.7)	(2.1)	(9.0)	(18.0)	(20.7)	(17.6)	(14.5)	(9.1)	(5.5)	(1.6)	(0.7)			(100.0)				
6～	7	26	77	293	605	721	607	536	383	179	55	11	1		3,501	249	3,750	155.15	9.37
(%)	(0.2)	(0.7)	(2.2)	(8.4)	(17.3)	(20.6)	(17.3)	(15.3)	(10.9)	(5.1)	(1.6)	(0.3)	(0.0)		(100.0)				
8～	12	13	102	383	1,015	1,694	1,637	1,448	1,102	632	192	40	7		8,277	470	8,747	157.53	9.07
(%)	(0.1)	(0.2)	(1.2)	(4.6)	(12.3)	(20.5)	(19.8)	(17.5)	(13.3)	(7.6)	(2.3)	(0.5)	(0.1)		(100.0)				
10～	9	10	36	189	658	1,435	1,824	2,067	1,746	1,108	412	94	7	3	9,598	549	10,147	160.76	8.87
(%)	(0.1)	(0.1)	(0.4)	(2.0)	(6.9)	(15.0)	(19.0)	(21.5)	(18.2)	(11.5)	(4.3)	(1.0)	(0.1)	(0.0)	(100.0)				
12～	1	1	4	38	157	493	893	1,403	1,500	1,202	431	140	19	1	6,283	355	6,638	164.51	7.92
(%)	(0.0)	(0.0)	(0.1)	(0.6)	(2.5)	(7.8)	(14.2)	(22.3)	(23.9)	(19.1)	(6.9)	(2.2)	(0.3)	(0.0)	(100.0)				
14～				8	14	66	191	499	747	730	303	80	21	2	2,661	159	2,820	167.79	6.97
(%)				(0.3)	(0.5)	(2.5)	(7.2)	(18.8)	(28.1)	(27.4)	(11.4)	(3.0)	(0.8)	(0.1)	(100.0)				
16～			1	1	3	13	44	155	281	276	156	60	7	1	998	50	1,048	169.40	6.76
(%)			(0.1)	(0.1)	(0.3)	(1.3)	(4.4)	(15.5)	(28.2)	(27.7)	(15.6)	(6.0)	(0.7)	(0.1)	(100.0)				
合計	34	56	240	1,002	2,618	4,621	5,359	6,246	5,857	4,176	1,562	431	62	7	32,271	1,947	34,218	160.73	9.52
(%)	(0.1)	(0.2)	(0.7)	(3.1)	(8.1)	(14.3)	(16.6)	(19.4)	(18.1)	(12.9)	(4.8)	(1.3)	(0.2)	(0.0)	(100.0)				
記載なし		1	6	5	14	43	50	56	43	35	9	2			264	489	753	160.23	9.17
(%)		(0.4)	(2.3)	(1.9)	(5.3)	(16.3)	(18.9)	(21.2)	(16.3)	(13.3)	(3.4)	(0.8)			(100.0)				
総計	34	57	246	1,007	2,632	4,664	5,409	6,302	5,900	4,211	1,571	433	62	7	32,535	2,436	34,971	160.72	9.52
(%)	(0.1)	(0.2)	(0.8)	(3.1)	(8.1)	(14.3)	(16.6)	(19.4)	(18.1)	(12.9)	(4.8)	(1.3)	(0.2)	(0.0)	(100.0)				
平均	8.56	8.05	8.38	8.54	9.01	9.62	10.22	10.92	11.54	12.09	12.50	12.90	13.45	13.24	10.74	10.44	10.72		
標準偏差	2.26	1.98	1.88	2.04	2.00	2.14	2.22	2.47	2.64	2.60	2.68	2.79	2.60	2.32	2.65	2.80	2.66		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査