

表 2406 身長(cm) PTH値(pg/mL)別（血液透析濾過患者 透析回数週3回）PTH測定法:whole-PTH

PTH値	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
20未満			3	10	25	39	35	37	40	30	13	1			233	16	249	159.67	9.79
(%)			(1.3)	(4.3)	(10.7)	(16.7)	(15.0)	(15.9)	(17.2)	(12.9)	(5.6)	(0.4)			(100.0)				
20～			3	7	30	49	53	51	49	31	9	2			284	17	301	159.47	8.67
(%)			(1.1)	(2.5)	(10.6)	(17.3)	(18.7)	(18.0)	(17.3)	(10.9)	(3.2)	(0.7)			(100.0)				
35～		1		10	24	42	48	45	40	30	12		1		253	21	274	159.56	9.15
(%)		(0.4)		(4.0)	(9.5)	(16.6)	(19.0)	(17.8)	(15.8)	(11.9)	(4.7)		(0.4)		(100.0)				
50～	1	1	1	15	38	51	60	85	87	48	19	5	1		412	32	444	160.75	9.34
(%)	(0.2)	(0.2)	(0.2)	(3.6)	(9.2)	(12.4)	(14.6)	(20.6)	(21.1)	(11.7)	(4.6)	(1.2)	(0.2)		(100.0)				
75～		2	5	7	31	51	61	79	69	50	14	5	2		376	15	391	160.78	9.36
(%)		(0.5)	(1.3)	(1.9)	(8.2)	(13.6)	(16.2)	(21.0)	(18.4)	(13.3)	(3.7)	(1.3)	(0.5)		(100.0)				
100～			2	6	25	35	53	64	53	48	14	7	1		308	23	331	161.84	9.17
(%)			(0.6)	(1.9)	(8.1)	(11.4)	(17.2)	(20.8)	(17.2)	(15.6)	(4.5)	(2.3)	(0.3)		(100.0)				
125～			3	6	11	19	35	47	52	35	4	3			215	19	234	161.92	8.67
(%)			(1.4)	(2.8)	(5.1)	(8.8)	(16.3)	(21.9)	(24.2)	(16.3)	(1.9)	(1.4)			(100.0)				
150～								2	2						4		4	163.25	3.95
(%)								(50.0)	(50.0)						(100.0)				
151～				11	16	31	31	32	40	24	19	1	1		206	9	215	161.17	9.99
(%)				(5.3)	(7.8)	(15.0)	(15.0)	(15.5)	(19.4)	(11.7)	(9.2)	(0.5)	(0.5)		(100.0)				
180～				5	9	12	27	22	18	26	8	1	2		130	7	137	162.21	9.64
(%)				(3.8)	(6.9)	(9.2)	(20.8)	(16.9)	(13.8)	(20.0)	(6.2)	(0.8)	(1.5)		(100.0)				
210～		1	1	4	7	8	12	15	22	12	11	1			94	4	98	162.35	10.40
(%)		(1.1)	(1.1)	(4.3)	(7.4)	(8.5)	(12.8)	(16.0)	(23.4)	(12.8)	(11.7)	(1.1)			(100.0)				
240～		1	1	3	3	11	12	13	17	14	5	1			81	7	88	161.67	9.68
(%)		(1.2)	(1.2)	(3.7)	(3.7)	(13.6)	(14.8)	(16.0)	(21.0)	(17.3)	(6.2)	(1.2)			(100.0)				
300～				1	5	5	6	3	6	9	2	1			38	4	42	162.16	10.63
(%)				(2.6)	(13.2)	(13.2)	(15.8)	(7.9)	(15.8)	(23.7)	(5.3)	(2.6)			(100.0)				
350～	1			1	2	4	4	4	9	1	1				27	2	29	159.22	10.54
(%)	(3.7)			(3.7)	(7.4)	(14.8)	(14.8)	(14.8)	(33.3)	(3.7)	(3.7)				(100.0)				
400～					2	6	3	3	3	3	2				22	2	24	160.73	9.60
(%)					(9.1)	(27.3)	(13.6)	(13.6)	(13.6)	(13.6)	(9.1)				(100.0)				
500～			1	1	6	7	8	6	7	5		1			42	3	45	158.52	9.47
(%)			(2.4)	(2.4)	(14.3)	(16.7)	(19.0)	(14.3)	(16.7)	(11.9)		(2.4)			(100.0)				
合計	2	6	20	87	234	370	448	508	514	366	133	29	8		2,725	181	2,906	160.79	9.40
(%)	(0.1)	(0.2)	(0.7)	(3.2)	(8.6)	(13.6)	(16.4)	(18.6)	(18.9)	(13.4)	(4.9)	(1.1)	(0.3)		(100.0)				
記載なし				1	1	5	4	3	3						17	14	31	156.59	7.24
(%)				(5.9)	(5.9)	(29.4)	(23.5)	(17.6)	(17.6)						(100.0)				
総計	2	6	20	88	235	375	452	511	517	366	133	29	8		2,742	195	2,937	160.77	9.39
(%)	(0.1)	(0.2)	(0.7)	(3.2)	(8.6)	(13.7)	(16.5)	(18.6)	(18.9)	(13.3)	(4.9)	(1.1)	(0.3)		(100.0)				
平均	222.00	125.33	110.45	110.06	110.34	108.61	109.54	107.06	114.09	118.23	117.61	128.00	115.88		111.79	111.76	111.79		
標準偏差	239.00	96.48	119.31	100.78	130.04	121.10	102.88	112.45	99.22	103.55	86.27	102.76	62.19		108.53	101.96	108.12		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査