

表 2426 身長(cm) クレアチニン濃度(mg/dL)別（腹膜透析患者）

クレアチニン濃度	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
2未満	1			1	1	5		3		1					12	5	17	151.00	17.35
(%)	(8.3)			(8.3)	(8.3)	(41.7)		(25.0)		(8.3)					(100.0)				
2～	8		5	19	16	24	22	20	14	7	1		1		137	17	154	150.85	19.67
(%)	(5.8)		(3.6)	(13.9)	(11.7)	(17.5)	(16.1)	(14.6)	(10.2)	(5.1)	(0.7)		(0.7)		(100.0)				
4～	21	3	14	45	61	90	97	99	56	40	12	6		1	545	30	575	154.52	15.72
(%)	(3.9)	(0.6)	(2.6)	(8.3)	(11.2)	(16.5)	(17.8)	(18.2)	(10.3)	(7.3)	(2.2)	(1.1)		(0.2)	(100.0)				
6～	10	2	20	60	114	184	208	184	169	86	27	5		1	1,070	63	1,133	157.39	11.62
(%)	(0.9)	(0.2)	(1.9)	(5.6)	(10.7)	(17.2)	(19.4)	(17.2)	(15.8)	(8.0)	(2.5)	(0.5)		(0.1)	(100.0)				
8～	10	2	12	40	112	211	219	239	215	155	60	10			1,285	64	1,349	159.59	10.75
(%)	(0.8)	(0.2)	(0.9)	(3.1)	(8.7)	(16.4)	(17.0)	(18.6)	(16.7)	(12.1)	(4.7)	(0.8)			(100.0)				
10～	3	1	4	17	57	132	169	258	202	150	65	9		2	1,069	46	1,115	161.92	8.88
(%)	(0.3)	(0.1)	(0.4)	(1.6)	(5.3)	(12.3)	(15.8)	(24.1)	(18.9)	(14.0)	(6.1)	(0.8)		(0.2)	(100.0)				
12～	1	3	1	5	21	54	122	194	181	154	58	13	4		811	37	848	164.20	8.30
(%)	(0.1)	(0.4)	(0.1)	(0.6)	(2.6)	(6.7)	(15.0)	(23.9)	(22.3)	(19.0)	(7.2)	(1.6)	(0.5)		(100.0)				
14～	2			2	5	12	43	97	123	109	48	18	2		461	27	488	166.86	8.14
(%)	(0.4)			(0.4)	(1.1)	(2.6)	(9.3)	(21.0)	(26.7)	(23.6)	(10.4)	(3.9)	(0.4)		(100.0)				
16～	1			1	3	11	18	36	69	75	36	17	4		271	24	295	168.31	9.01
(%)	(0.4)			(0.4)	(1.1)	(4.1)	(6.6)	(13.3)	(25.5)	(27.7)	(13.3)	(6.3)	(1.5)		(100.0)				
合計	57	11	56	190	390	723	898	1,130	1,029	777	307	78	11	4	5,661	313	5,974	160.57	11.65
(%)	(1.0)	(0.2)	(1.0)	(3.4)	(6.9)	(12.8)	(15.9)	(20.0)	(18.2)	(13.7)	(5.4)	(1.4)	(0.2)	(0.1)	(100.0)				
記載なし	5		6	37	80	144	151	200	183	127	59	9	2		1,003	1,964	2,967	160.50	11.26
(%)	(0.5)		(0.6)	(3.7)	(8.0)	(14.4)	(15.1)	(19.9)	(18.2)	(12.7)	(5.9)	(0.9)	(0.2)		(100.0)				
総計	62	11	62	227	470	867	1,049	1,330	1,212	904	366	87	13	4	6,664	2,277	8,941	160.56	11.59
(%)	(0.9)	(0.2)	(0.9)	(3.4)	(7.1)	(13.0)	(15.7)	(20.0)	(18.2)	(13.6)	(5.5)	(1.3)	(0.2)	(0.1)	(100.0)				
平均	6.72	8.61	6.84	7.12	8.12	8.60	9.32	10.07	10.74	11.36	11.79	12.83	14.09	8.48	9.90	9.65	9.89		
標準偏差	3.30	3.21	2.23	2.64	2.62	2.83	3.25	3.19	3.53	3.58	3.49	4.02	4.30	3.45	3.50	3.97	3.53		

数値下のかっこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2014年調査