

表 2431 身長(cm) PTH値(pg/mL)別（腹膜透析患者）PTH測定法:intact-PTH

PTH値	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
20未満	3	1		5	7	14	13	29	24	15	5	1			117	6	123	159.52	14.87
(%)	(2.6)	(0.9)		(4.3)	(6.0)	(12.0)	(11.1)	(24.8)	(20.5)	(12.8)	(4.3)	(0.9)			(100.0)				
20～	7		3	10	25	30	35	49	56	37	10	5		1	268	13	281	159.67	13.58
(%)	(2.6)		(1.1)	(3.7)	(9.3)	(11.2)	(13.1)	(18.3)	(20.9)	(13.8)	(3.7)	(1.9)		(0.4)	(100.0)				
40～	1	1	3	9	17	28	40	50	45	39	18	2			253	15	268	161.17	11.02
(%)	(0.4)	(0.4)	(1.2)	(3.6)	(6.7)	(11.1)	(15.8)	(19.8)	(17.8)	(15.4)	(7.1)	(0.8)			(100.0)				
60～	3		4	15	22	43	63	54	55	29	9			1	298	17	315	158.53	11.82
(%)	(1.0)		(1.3)	(5.0)	(7.4)	(14.4)	(21.1)	(18.1)	(18.5)	(9.7)	(3.0)			(0.3)	(100.0)				
80～	4		2	6	13	42	48	62	62	36	16	3			294	15	309	160.87	11.08
(%)	(1.4)		(0.7)	(2.0)	(4.4)	(14.3)	(16.3)	(21.1)	(21.1)	(12.2)	(5.4)	(1.0)			(100.0)				
100～	3		6	13	29	41	54	67	39	33	13	6			304	12	316	158.81	12.01
(%)	(1.0)		(2.0)	(4.3)	(9.5)	(13.5)	(17.8)	(22.0)	(12.8)	(10.9)	(4.3)	(2.0)			(100.0)				
120～	2		3	11	17	39	43	56	57	34	21	2	1		286	19	305	161.07	10.19
(%)	(0.7)		(1.0)	(3.8)	(5.9)	(13.6)	(15.0)	(19.6)	(19.9)	(11.9)	(7.3)	(0.7)	(0.3)		(100.0)				
140～	1			9	20	42	28	64	51	37	10	5			267	18	285	161.05	10.02
(%)	(0.4)			(3.4)	(7.5)	(15.7)	(10.5)	(24.0)	(19.1)	(13.9)	(3.7)	(1.9)			(100.0)				
160～	3	1		7	15	34	40	57	38	36	20	4	1		259	17	276	160.65	12.97
(%)	(1.2)	(0.4)		(2.7)	(5.8)	(13.1)	(15.4)	(22.0)	(14.7)	(13.9)	(7.7)	(1.5)	(0.4)		(100.0)				
180～	4	2	5	18	33	87	96	126	121	88	36	11	2	2	631	23	654	161.42	11.26
(%)	(0.6)	(0.3)	(0.8)	(2.9)	(5.2)	(13.8)	(15.2)	(20.0)	(19.2)	(13.9)	(5.7)	(1.7)	(0.3)	(0.3)	(100.0)				
240～					1	2	3	3		1	3				10		10	165.70	9.72
(%)					(10.0)	(20.0)	(30.0)	(30.0)		(10.0)	(30.0)				(100.0)				
241～	4	1	1	8	27	53	82	83	77	50	17	8	1		412	19	431	160.75	10.53
(%)	(1.0)	(0.2)	(0.2)	(1.9)	(6.6)	(12.9)	(19.9)	(20.1)	(18.7)	(12.1)	(4.1)	(1.9)	(0.2)		(100.0)				
300～		1	5	6	18	36	52	60	58	51	21	3			311	20	331	161.96	9.33
(%)		(0.3)	(1.6)	(1.9)	(5.8)	(11.6)	(16.7)	(19.3)	(18.6)	(16.4)	(6.8)	(1.0)			(100.0)				
360～	3		3	4	12	18	28	37	43	31	8	2			189	11	200	160.74	13.38
(%)	(1.6)		(1.6)	(2.1)	(6.3)	(9.5)	(14.8)	(19.6)	(22.8)	(16.4)	(4.2)	(1.1)			(100.0)				
420～	2		3	7	9	18	17	25	21	23	3	1			129	10	139	159.28	11.77
(%)	(1.6)		(2.3)	(5.4)	(7.0)	(14.0)	(13.2)	(19.4)	(16.3)	(17.8)	(2.3)	(0.8)			(100.0)				
480～				3	5	4	3	3	9	8	5	1			41	2	43	162.93	11.21
(%)				(7.3)	(12.2)	(9.8)	(7.3)	(7.3)	(22.0)	(19.5)	(12.2)	(2.4)			(100.0)				
500～		1		4	2	5	10	11	5	8	6	1			53	1	54	161.36	10.98
(%)		(1.9)		(7.5)	(3.8)	(9.4)	(18.9)	(20.8)	(9.4)	(15.1)	(11.3)	(1.9)			(100.0)				
540～				1	4	5	15	14	7	12	5	1	1		65	8	73	162.82	9.25
(%)				(1.5)	(6.2)	(7.7)	(23.1)	(21.5)	(10.8)	(18.5)	(7.7)	(1.5)	(1.5)		(100.0)				
600～	3		2	2	15	9	22	18	9	14	5	1			100	5	105	158.19	12.04
(%)	(3.0)		(2.0)	(2.0)	(15.0)	(9.0)	(22.0)	(18.0)	(9.0)	(14.0)	(5.0)	(1.0)			(100.0)				
800～	5		2	5	7	16	20	14	15	12	2	2			100	10	110	155.70	19.10
(%)	(5.0)		(2.0)	(5.0)	(7.0)	(16.0)	(20.0)	(14.0)	(15.0)	(12.0)	(2.0)	(2.0)			(100.0)				
合計	48	7	46	143	297	565	711	882	792	594	233	59	6	4	4,387	241	4,628	160.46	11.76
(%)	(1.1)	(0.2)	(1.0)	(3.3)	(6.8)	(12.9)	(16.2)	(20.1)	(18.1)	(13.5)	(5.3)	(1.3)	(0.1)	(0.1)	(100.0)				
記載なし		1	1	11	14	28	52	57	61	40	15	5			285	91	376	161.97	9.20
(%)		(0.4)	(0.4)	(3.9)	(4.9)	(9.8)	(18.2)	(20.0)	(21.4)	(14.0)	(5.3)	(1.8)			(100.0)				
総計	48	8	47	154	311	593	763	939	853	634	248	64	6	4	4,672	332	5,004	160.56	11.62
(%)	(1.0)	(0.2)	(1.0)	(3.3)	(6.7)	(12.7)	(16.3)	(20.1)	(18.3)	(13.6)	(5.3)	(1.4)	(0.1)	(0.1)	(100.0)				
平均	306.83	174.29	269.85	247.50	238.68	216.92	234.14	209.94	209.08	229.71	219.21	240.20	262.50	137.50	222.99	240.00	223.88		
標準偏差	400.77	111.71	302.19	393.27	269.31	217.42	257.18	193.91	204.22	201.43	184.55	225.59	164.85	100.63	229.54	224.90	229.31		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査