

表 2443 透析後体重(kg) 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	26	293	6,829	20,494	21,926	11,712	3,768	1,135	362	122	59	34	66,760	1,817	68,577	53.45	11.87
(%)	(0.0)	(0.4)	(10.2)	(30.7)	(32.8)	(17.5)	(5.6)	(1.7)	(0.5)	(0.2)	(0.1)	(0.1)	(100.0)				
慢性腎盂腎炎		20	339	692	643	303	77	32	8	3	1	1	2,119	46	2,165	51.08	11.81
(%)		(0.9)	(16.0)	(32.7)	(30.3)	(14.3)	(3.6)	(1.5)	(0.4)	(0.1)	(0.0)	(0.0)	(100.0)				
急速進行性糸球体腎炎	1	15	292	616	500	261	67	17	6				1,775	59	1,834	50.48	11.18
(%)	(0.1)	(0.8)	(16.5)	(34.7)	(28.2)	(14.7)	(3.8)	(1.0)	(0.3)				(100.0)				
妊娠腎/妊娠中毒症		7	176	459	284	78	25	11	2	1			1,043	21	1,064	48.68	10.13
(%)		(0.7)	(16.9)	(44.0)	(27.2)	(7.5)	(2.4)	(1.1)	(0.2)	(0.1)			(100.0)				
その他の分類不能の腎炎	1	8	111	275	288	157	53	18	4	4		2	921	28	949	53.27	13.20
(%)	(0.1)	(0.9)	(12.1)	(29.9)	(31.3)	(17.0)	(5.8)	(2.0)	(0.4)	(0.4)		(0.2)	(100.0)				
多発性嚢胞腎	3	12	481	2,180	2,633	1,732	677	193	59	16	7	6	7,999	199	8,198	56.01	12.00
(%)	(0.0)	(0.2)	(6.0)	(27.3)	(32.9)	(21.7)	(8.5)	(2.4)	(0.7)	(0.2)	(0.1)	(0.1)	(100.0)				
腎硬化症	5	109	2,343	6,351	6,722	3,550	1,216	387	149	49	22	12	20,915	545	21,460	53.39	12.31
(%)	(0.0)	(0.5)	(11.2)	(30.4)	(32.1)	(17.0)	(5.8)	(1.9)	(0.7)	(0.2)	(0.1)	(0.1)	(100.0)				
悪性高血圧	1	8	148	465	524	358	186	80	39	7	3	4	1,823	58	1,881	57.25	14.63
(%)	(0.1)	(0.4)	(8.1)	(25.5)	(28.7)	(19.6)	(10.2)	(4.4)	(2.1)	(0.4)	(0.2)	(0.2)	(100.0)				
糖尿病性腎症	39	126	4,503	18,799	28,704	20,754	9,319	3,639	1,370	486	173	111	88,023	2,484	90,507	58.52	13.41
(%)	(0.0)	(0.1)	(5.1)	(21.4)	(32.6)	(23.6)	(10.6)	(4.1)	(1.6)	(0.6)	(0.2)	(0.1)	(100.0)				
SLE腎炎		10	269	618	387	167	42	13	6				1,512	38	1,550	49.18	10.72
(%)		(0.7)	(17.8)	(40.9)	(25.6)	(11.0)	(2.8)	(0.9)	(0.4)				(100.0)				
アミロイド腎	6	83	100	84	29	8	1						311	14	325	47.51	10.90
(%)		(1.9)	(26.7)	(32.2)	(27.0)	(9.3)	(2.6)	(0.3)					(100.0)				
痛風腎	1	48	165	292	205	68	18	8	2		1		808	20	828	57.16	11.49
(%)		(0.1)	(5.9)	(20.4)	(36.1)	(25.4)	(8.4)	(2.2)	(1.0)	(0.2)	(0.1)		(100.0)				
先天性代謝異常による腎不全		1	24	54	53	25	11	5	1				174	3	177	52.75	12.89
(%)		(0.6)	(13.8)	(31.0)	(30.5)	(14.4)	(6.3)	(2.9)	(0.6)				(100.0)				
腎・尿路結核		1	32	47	52	21							153	5	158	48.81	9.99
(%)		(0.7)	(20.9)	(30.7)	(34.0)	(13.7)							(100.0)				
腎・尿路結石	1	3	48	129	130	77	27	1	3	2	1		422	8	430	53.00	12.57
(%)	(0.2)	(0.7)	(11.4)	(30.6)	(30.8)	(18.2)	(6.4)	(0.2)	(0.7)	(0.5)	(0.2)		(100.0)				
腎・尿路腫瘍		1	50	179	242	138	44	12	3		1		670	16	686	54.98	11.33
(%)		(0.1)	(7.5)	(26.7)	(36.1)	(20.6)	(6.6)	(1.8)	(0.4)		(0.1)		(100.0)				
閉塞性尿路障害		3	70	146	164	82	26	9	1	1			502	15	517	52.45	11.47
(%)		(0.6)	(13.9)	(29.1)	(32.7)	(16.3)	(5.2)	(1.8)	(0.2)	(0.2)			(100.0)				
骨髄腫		2	22	65	61	21	10	2					183	7	190	51.07	10.10
(%)		(1.1)	(12.0)	(35.5)	(33.3)	(11.5)	(5.5)	(1.1)					(100.0)				
腎形成不全	1	5	53	121	117	72	25	12	3			1	410	9	419	53.23	13.41
(%)	(0.2)	(1.2)	(12.9)	(29.5)	(28.5)	(17.6)	(6.1)	(2.9)	(0.7)			(0.2)	(100.0)				
不明	5	126	2,423	5,893	6,118	3,216	1,102	368	123	45	26	12	19,457	668	20,125	53.07	12.48
(%)	(0.0)	(0.6)	(12.5)	(30.3)	(31.4)	(16.5)	(5.7)	(1.9)	(0.6)	(0.2)	(0.1)	(0.1)	(100.0)				
再導入		9	139	406	442	268	86	24	10	1			1,385	32	1,417	53.85	11.68
(%)		(0.6)	(10.0)	(29.3)	(31.9)	(19.4)	(6.2)	(1.7)	(0.7)	(0.1)			(100.0)				
その他	1	36	634	1,555	1,743	856	347	104	38	16	5	6	5,341	173	5,514	53.47	12.68
(%)	(0.0)	(0.7)	(11.9)	(29.1)	(32.6)	(16.0)	(6.5)	(1.9)	(0.7)	(0.3)	(0.1)	(0.1)	(100.0)				
合計	84	802	19,117	59,809	72,109	44,082	17,184	6,081	2,195	755	299	189	222,706	6,265	228,971	55.45	12.90
(%)	(0.0)	(0.4)	(8.6)	(26.9)	(32.4)	(19.8)	(7.7)	(2.7)	(1.0)	(0.3)	(0.1)	(0.1)	(100.0)				
記載なし																	
(%)																	
総計	84	802	19,117	59,809	72,109	44,082	17,184	6,081	2,195	755	299	189	222,706	6,265	228,971	55.45	12.90
(%)	(0.0)	(0.4)	(8.6)	(26.9)	(32.4)	(19.8)	(7.7)	(2.7)	(1.0)	(0.3)	(0.1)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査