

表 2456 透析後体重(kg) 腎移植の回数別（施設血液透析患者 透析回数週3回）

腎移植の回数	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
なし	67	642	15,333	47,673	57,277	35,070	13,681	4,818	1,735	591	227	151	177,265	3,506	180,771	55.41	12.90
(%)	(0.0)	(0.4)	(8.6)	(26.9)	(32.3)	(19.8)	(7.7)	(2.7)	(1.0)	(0.3)	(0.1)	(0.1)	(100.0)				
1回		8	141	535	601	306	89	37	15	4	2		1,738	26	1,764	54.04	11.75
(%)		(0.5)	(8.1)	(30.8)	(34.6)	(17.6)	(5.1)	(2.1)	(0.9)	(0.2)	(0.1)		(100.0)				
2回		1	10	29	24	18	3			1			86	2	88	52.33	12.12
(%)		(1.2)	(11.6)	(33.7)	(27.9)	(20.9)	(3.5)			(1.2)			(100.0)				
3回			1	4									5		5	42.40	4.73
(%)			(20.0)	(80.0)									(100.0)				
4回以上																	
(%)																	
合計	67	651	15,485	48,241	57,902	35,394	13,773	4,855	1,750	596	229	151	179,094	3,534	182,628	55.40	12.89
(%)	(0.0)	(0.4)	(8.6)	(26.9)	(32.3)	(19.8)	(7.7)	(2.7)	(1.0)	(0.3)	(0.1)	(0.1)	(100.0)				
不明	1	3	74	185	232	109	40	12	5	2	2		665	22	687	53.67	12.41
(%)	(0.2)	(0.5)	(11.1)	(27.8)	(34.9)	(16.4)	(6.0)	(1.8)	(0.8)	(0.3)	(0.3)		(100.0)				
記載なし	16	148	3,558	11,383	13,975	8,579	3,371	1,214	440	157	68	38	42,947	2,709	45,656	55.67	12.95
(%)	(0.0)	(0.3)	(8.3)	(26.5)	(32.5)	(20.0)	(7.8)	(2.8)	(1.0)	(0.4)	(0.2)	(0.1)	(100.0)				
総計	84	802	19,117	59,809	72,109	44,082	17,184	6,081	2,195	755	299	189	222,706	6,265	228,971	55.45	12.90
(%)	(0.0)	(0.4)	(8.6)	(26.9)	(32.4)	(19.8)	(7.7)	(2.7)	(1.0)	(0.3)	(0.1)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査