

表 2459 透析後体重(kg) 身長(cm)別（施設血液透析患者 透析回数週3回）

身長	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
130未満	1	24	93	69	27	10	8	1	1				234	3	237	41.85	12.46
(%)	(0.4)	(10.3)	(39.7)	(29.5)	(11.5)	(4.3)	(3.4)	(0.4)	(0.4)				(100.0)				
130～		65	320	134	33	6	2						560	8	568	37.69	7.60
(%)		(11.6)	(57.1)	(23.9)	(5.9)	(1.1)	(0.4)						(100.0)				
135～	1	106	1,159	805	163	30	2		1			1	2,268	37	2,305	40.01	8.84
(%)	(0.0)	(4.7)	(51.1)	(35.5)	(7.2)	(1.3)	(0.1)		(0.0)			(0.0)	(100.0)				
140～	3	221	3,240	3,550	907	212	33	8	3	2			8,179	104	8,283	42.29	7.97
(%)	(0.0)	(2.7)	(39.6)	(43.4)	(11.1)	(2.6)	(0.4)	(0.1)	(0.0)	(0.0)			(100.0)				
145～	6	152	5,029	8,867	3,489	803	147	44	4	1	1		18,543	276	18,819	45.13	8.38
(%)	(0.0)	(0.8)	(27.1)	(47.8)	(18.8)	(4.3)	(0.8)	(0.2)	(0.0)	(0.0)	(0.0)		(100.0)				
150～	14	87	4,466	13,782	8,267	2,286	550	122	31	5	5		29,615	399	30,014	48.46	9.07
(%)	(0.0)	(0.3)	(15.1)	(46.5)	(27.9)	(7.7)	(1.9)	(0.4)	(0.1)	(0.0)	(0.0)		(100.0)				
155～	16	44	2,110	12,980	13,379	4,498	986	270	75	16	11	2	34,387	470	34,857	52.11	9.28
(%)	(0.0)	(0.1)	(6.1)	(37.7)	(38.9)	(13.1)	(2.9)	(0.8)	(0.2)	(0.0)	(0.0)	(0.0)	(100.0)				
160～	10	7	744	9,340	18,707	9,446	2,592	637	159	50	11	8	41,711	547	42,258	56.52	9.58
(%)	(0.0)	(0.0)	(1.8)	(22.4)	(44.8)	(22.6)	(6.2)	(1.5)	(0.4)	(0.1)	(0.0)	(0.0)	(100.0)				
165～	17	4	194	4,349	14,217	12,010	4,443	1,303	404	97	34	24	37,096	480	37,576	60.98	10.63
(%)	(0.0)	(0.0)	(0.5)	(11.7)	(38.3)	(32.4)	(12.0)	(3.5)	(1.1)	(0.3)	(0.1)	(0.1)	(100.0)				
170～	4	2	60	1,245	6,420	8,665	4,482	1,714	609	197	76	43	23,517	310	23,827	65.83	12.00
(%)	(0.0)	(0.0)	(0.3)	(5.3)	(27.3)	(36.8)	(19.1)	(7.3)	(2.6)	(0.8)	(0.3)	(0.2)	(100.0)				
175～	1	1	11	186	1,413	2,761	2,204	1,109	466	180	70	47	8,449	100	8,549	71.29	13.70
(%)	(0.0)	(0.0)	(0.1)	(2.2)	(16.7)	(32.7)	(26.1)	(13.1)	(5.5)	(2.1)	(0.8)	(0.6)	(100.0)				
180～	1	1	1	19	191	549	611	418	242	105	55	38	2,231	23	2,254	77.84	15.80
(%)	(0.0)	(0.0)	(0.0)	(0.9)	(8.6)	(24.6)	(27.4)	(18.7)	(10.8)	(4.7)	(2.5)	(1.7)	(100.0)				
185～			1	1	13	60	88	80	47	37	9	11	347	4	351	83.19	16.69
(%)			(0.3)	(0.3)	(3.7)	(17.3)	(25.4)	(23.1)	(13.5)	(10.7)	(2.6)	(3.2)	(100.0)				
190～			1	2	4	6	7	7	7	12	7	3	56	1	57	88.89	23.73
(%)			(1.8)	(3.6)	(7.1)	(10.7)	(12.5)	(12.5)	(12.5)	(21.4)	(12.5)	(5.4)	(100.0)				
合計	74	714	17,429	55,329	67,230	41,342	16,155	5,713	2,049	702	279	177	207,193	2,762	209,955	55.55	12.89
(%)	(0.0)	(0.3)	(8.4)	(26.7)	(32.4)	(20.0)	(7.8)	(2.8)	(1.0)	(0.3)	(0.1)	(0.1)	(100.0)				
記載なし	10	88	1,688	4,480	4,879	2,740	1,029	368	146	53	20	12	15,513	3,503	19,016	54.08	12.97
(%)	(0.1)	(0.6)	(10.9)	(28.9)	(31.5)	(17.7)	(6.6)	(2.4)	(0.9)	(0.3)	(0.1)	(0.1)	(100.0)				
総計	84	802	19,117	59,809	72,109	44,082	17,184	6,081	2,195	755	299	189	222,706	6,265	228,971	55.45	12.90
(%)	(0.0)	(0.4)	(8.6)	(26.9)	(32.4)	(19.8)	(7.7)	(2.7)	(1.0)	(0.3)	(0.1)	(0.1)	(100.0)				
平均	156.97	143.12	148.21	154.65	160.88	165.19	168.10	170.19	171.88	173.80	174.08	175.07	159.95	159.45	159.94		
標準偏差	14.65	8.44	7.15	7.67	7.47	7.30	7.32	7.31	7.27	7.64	7.81	7.57	9.59	9.56	9.59		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査