

表 2460 透析後体重(kg) 身長(cm)別 (施設血液透析患者 男性のみ 透析回数週3回)

身長	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
130未満	1	4	18	33	15	6	8	1					86	1	87	47.78	14.16
(%)	(1.2)	(4.7)	(20.9)	(38.4)	(17.4)	(7.0)	(9.3)	(1.2)					(100.0)				
130～		3	12	12	13	3	1						44	1	45	46.00	10.76
(%)		(6.8)	(27.3)	(27.3)	(29.5)	(6.8)	(2.3)						(100.0)				
135～		2	38	24	20	9						1	94		94	47.14	26.22
(%)		(2.1)	(40.4)	(25.5)	(21.3)	(9.6)						(1.1)	(100.0)				
140～	1	6	117	183	55	21	3	2	2				390	7	397	44.55	9.70
(%)	(0.3)	(1.5)	(30.0)	(46.9)	(14.1)	(5.4)	(0.8)	(0.5)	(0.5)				(100.0)				
145～		4	298	870	360	63	14	3					1,612	22	1,634	46.27	7.61
(%)		(0.2)	(18.5)	(54.0)	(22.3)	(3.9)	(0.9)	(0.2)					(100.0)				
150～	1	3	618	3,357	2,554	506	62	11	2	2			7,116	97	7,213	49.39	7.52
(%)	(0.0)	(0.0)	(8.7)	(47.2)	(35.9)	(7.1)	(0.9)	(0.2)	(0.0)	(0.0)			(100.0)				
155～	8	3	661	6,577	8,482	2,652	434	70	23	1	3		18,914	248	19,162	52.73	8.13
(%)	(0.0)	(0.0)	(3.5)	(34.8)	(44.8)	(14.0)	(2.3)	(0.4)	(0.1)	(0.0)	(0.0)		(100.0)				
160～	10	1	467	7,321	16,366	8,382	2,138	470	95	29	5	4	35,288	456	35,744	56.72	9.09
(%)	(0.0)	(0.0)	(1.3)	(20.7)	(46.4)	(23.8)	(6.1)	(1.3)	(0.3)	(0.1)	(0.0)	(0.0)	(100.0)				
165～	16	4	154	4,029	13,667	11,630	4,272	1,229	368	89	33	21	35,512	459	35,971	61.03	10.51
(%)	(0.0)	(0.0)	(0.4)	(11.3)	(38.5)	(32.7)	(12.0)	(3.5)	(1.0)	(0.3)	(0.1)	(0.1)	(100.0)				
170～	4	2	53	1,206	6,337	8,571	4,436	1,695	600	192	76	43	23,215	305	23,520	65.86	11.98
(%)	(0.0)	(0.0)	(0.2)	(5.2)	(27.3)	(36.9)	(19.1)	(7.3)	(2.6)	(0.8)	(0.3)	(0.2)	(100.0)				
175～	1	1	7	178	1,399	2,742	2,186	1,104	464	179	69	46	8,376	99	8,475	71.34	13.65
(%)	(0.0)	(0.0)	(0.1)	(2.1)	(16.7)	(32.7)	(26.1)	(13.2)	(5.5)	(2.1)	(0.8)	(0.5)	(100.0)				
180～	1	1	1	18	190	547	608	417	242	105	55	38	2,223	23	2,246	77.88	15.80
(%)	(0.0)	(0.0)	(0.0)	(0.8)	(8.5)	(24.6)	(27.4)	(18.8)	(10.9)	(4.7)	(2.5)	(1.7)	(100.0)				
185～			1	1	13	60	86	80	46	37	9	11	344	4	348	83.20	16.76
(%)			(0.3)	(0.3)	(3.8)	(17.4)	(25.0)	(23.3)	(13.4)	(10.8)	(2.6)	(3.2)	(100.0)				
190～			1	2	4	6	7	7	7	12	7	3	56		56	88.89	23.73
(%)			(1.8)	(3.6)	(7.1)	(10.7)	(12.5)	(12.5)	(12.5)	(21.4)	(12.5)	(5.4)	(100.0)				
合計	43	34	2,446	23,811	49,475	35,198	14,255	5,089	1,849	646	257	167	133,270	1,722	134,992	59.68	12.17
(%)	(0.0)	(0.0)	(1.8)	(17.9)	(37.1)	(26.4)	(10.7)	(3.8)	(1.4)	(0.5)	(0.2)	(0.1)	(100.0)				
記載なし	6	6	272	2,100	3,652	2,318	892	329	125	48	16	11	9,775	2,266	12,041	58.27	12.24
(%)	(0.1)	(0.1)	(2.8)	(21.5)	(37.4)	(23.7)	(9.1)	(3.4)	(1.3)	(0.5)	(0.2)	(0.1)	(100.0)				
総計	49	40	2,718	25,911	53,127	37,516	15,147	5,418	1,974	694	273	178	143,045	3,988	147,033	59.58	12.18
(%)	(0.0)	(0.0)	(1.9)	(18.1)	(37.1)	(26.2)	(10.6)	(3.8)	(1.4)	(0.5)	(0.2)	(0.1)	(100.0)				
平均	161.21	148.38	154.78	159.67	163.48	166.90	169.54	171.63	173.13	174.93	175.41	175.69	164.72	164.42	164.72		
標準偏差	17.48	16.95	8.07	6.55	6.07	5.96	6.11	6.04	6.12	6.61	6.35	7.19	7.26	7.11	7.26		

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査