

表 2462 透析後体重(kg) 体重減少量(kg)別（施設血液透析患者 透析回数週3回 透析歴2年以上）

体重減少量	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
0.0未満		8	121	227	237	120	32	5	2	3			755		755	51.32	11.81
(%)		(1.1)	(16.0)	(30.1)	(31.4)	(15.9)	(4.2)	(0.7)	(0.3)	(0.4)			(100.0)				
0.0～	2	144	1,660	3,096	2,278	949	251	58	5	4		3	8,450		8,450	48.86	10.91
(%)	(0.0)	(1.7)	(19.6)	(36.6)	(27.0)	(11.2)	(3.0)	(0.7)	(0.1)	(0.0)		(0.0)	(100.0)				
1.0～		358	7,079	15,530	12,000	4,717	1,059	220	47	8	6	1	41,025		41,025	49.34	10.25
(%)		(0.9)	(17.3)	(37.9)	(29.3)	(11.5)	(2.6)	(0.5)	(0.1)	(0.0)	(0.0)	(0.0)	(100.0)				
2.0～		126	5,737	22,292	26,257	13,749	3,893	865	185	39	12	6	73,161		73,161	53.67	10.52
(%)		(0.2)	(7.8)	(30.5)	(35.9)	(18.8)	(5.3)	(1.2)	(0.3)	(0.1)	(0.0)	(0.0)	(100.0)				
3.0～		6	704	5,736	13,016	11,146	5,162	1,653	520	120	33	14	38,110		38,110	60.76	11.75
(%)		(0.0)	(1.8)	(15.1)	(34.2)	(29.2)	(13.5)	(4.3)	(1.4)	(0.3)	(0.1)	(0.0)	(100.0)				
4.0～		1	33	496	1,968	2,803	2,106	1,211	545	178	68	36	9,445		9,445	69.64	14.36
(%)		(0.0)	(0.3)	(5.3)	(20.8)	(29.7)	(22.3)	(12.8)	(5.8)	(1.9)	(0.7)	(0.4)	(100.0)				
5.0～		1	8	72	249	384	450	344	228	127	55	40	1,958		1,958	77.34	18.16
(%)		(0.1)	(0.4)	(3.7)	(12.7)	(19.6)	(23.0)	(17.6)	(11.6)	(6.5)	(2.8)	(2.0)	(100.0)				
6.0～			2	9	26	57	63	74	63	47	29	25	395		395	86.67	21.60
(%)			(0.5)	(2.3)	(6.6)	(14.4)	(15.9)	(18.7)	(15.9)	(11.9)	(7.3)	(6.3)	(100.0)				
7.0～			2	11	11	13	10	6	13	9	9	11	95		95	83.84	28.92
(%)			(2.1)	(11.6)	(11.6)	(13.7)	(10.5)	(6.3)	(13.7)	(9.5)	(9.5)	(11.6)	(100.0)				
8.0～	4	3	15	48	45	17	8	6	4	5	3	2	160		160	55.96	21.71
(%)	(2.5)	(1.9)	(9.4)	(30.0)	(28.1)	(10.6)	(5.0)	(3.8)	(2.5)	(3.1)	(1.9)	(1.3)	(100.0)				
合計	6	647	15,361	47,517	56,087	33,955	13,034	4,442	1,612	540	215	138	173,554		173,554	55.19	12.75
(%)	(0.0)	(0.4)	(8.9)	(27.4)	(32.3)	(19.6)	(7.5)	(2.6)	(0.9)	(0.3)	(0.1)	(0.1)	(100.0)				
記載なし	56	9	11	16	19	11	7	9	3				141	4,514	4,655	36.98	25.90
(%)	(39.7)	(6.4)	(7.8)	(11.3)	(13.5)	(7.8)	(5.0)	(6.4)	(2.1)				(100.0)				
総計	62	656	15,372	47,533	56,106	33,966	13,041	4,451	1,615	540	215	138	173,695	4,514	178,209	55.17	12.78
(%)	(0.0)	(0.4)	(8.8)	(27.4)	(32.3)	(19.6)	(7.5)	(2.6)	(0.9)	(0.3)	(0.1)	(0.1)	(100.0)				
平均	13.97	1.47	1.81	2.14	2.46	2.79	3.18	3.60	4.06	4.49	4.82	5.13	2.49		2.49		
標準偏差	11.78	0.96	0.81	0.90	0.92	0.98	1.04	1.11	1.14	1.34	1.36	1.49	1.05		1.05		

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査