

表 2463 透析後体重(kg) 体重減少量(kg)別（施設血液透析患者 男性のみ 透析回数週3回 透析歴2年以上）

体重減少量	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
0.0未満		1	16	125	174	104	28	5	2	2			457		457	55.64	10.63
(%)		(0.2)	(3.5)	(27.4)	(38.1)	(22.8)	(6.1)	(1.1)	(0.4)	(0.4)			(100.0)				
0.0～	2	6	313	1,417	1,683	805	212	52	4	4		2	4,500		4,500	53.62	10.20
(%)	(0.0)	(0.1)	(7.0)	(31.5)	(37.4)	(17.9)	(4.7)	(1.2)	(0.1)	(0.1)		(0.0)	(100.0)				
1.0～		6	908	6,071	8,253	3,852	917	188	37	6	6	1	20,245		20,245	54.25	9.52
(%)		(0.0)	(4.5)	(30.0)	(40.8)	(19.0)	(4.5)	(0.9)	(0.2)	(0.0)	(0.0)	(0.0)	(100.0)				
2.0～		5	753	9,249	18,509	11,297	3,292	721	162	32	10	4	44,034		44,034	57.19	9.67
(%)		(0.0)	(1.7)	(21.0)	(42.0)	(25.7)	(7.5)	(1.6)	(0.4)	(0.1)	(0.0)	(0.0)	(100.0)				
3.0～		1	125	2,867	10,309	9,717	4,592	1,460	445	111	27	13	29,667		29,667	62.58	11.07
(%)		(0.0)	(0.4)	(9.7)	(34.7)	(32.8)	(15.5)	(4.9)	(1.5)	(0.4)	(0.1)	(0.0)	(100.0)				
4.0～		1	9	284	1,664	2,547	1,935	1,111	505	167	63	33	8,319		8,319	70.54	13.98
(%)		(0.0)	(0.1)	(3.4)	(20.0)	(30.6)	(23.3)	(13.4)	(6.1)	(2.0)	(0.8)	(0.4)	(100.0)				
5.0～			1	37	224	360	428	323	221	125	52	38	1,809		1,809	78.36	17.62
(%)			(0.1)	(2.0)	(12.4)	(19.9)	(23.7)	(17.9)	(12.2)	(6.9)	(2.9)	(2.1)	(100.0)				
6.0～			1	4	23	56	60	72	61	46	28	25	376		376	87.67	21.01
(%)			(0.3)	(1.1)	(6.1)	(14.9)	(16.0)	(19.1)	(16.2)	(12.2)	(7.4)	(6.6)	(100.0)				
7.0～				6	9	12	10	6	12	8	9	11	83		83	87.99	27.60
(%)				(7.2)	(10.8)	(14.5)	(12.0)	(7.2)	(14.5)	(9.6)	(10.8)	(13.3)	(100.0)				
8.0～	1		5	35	36	13	6	6	4	5	3	2	116		116	60.55	21.96
(%)	(0.9)		(4.3)	(30.2)	(31.0)	(11.2)	(5.2)	(5.2)	(3.4)	(4.3)	(2.6)	(1.7)	(100.0)				
合計	3	20	2,131	20,095	40,884	28,763	11,480	3,944	1,453	506	198	129	109,606		109,606	59.45	12.02
(%)	(0.0)	(0.0)	(1.9)	(18.3)	(37.3)	(26.2)	(10.5)	(3.6)	(1.3)	(0.5)	(0.2)	(0.1)	(100.0)				
記載なし	30	7	8	10	15	9	4	6	2				91	2,818	2,909	39.65	25.37
(%)	(33.0)	(7.7)	(8.8)	(11.0)	(16.5)	(9.9)	(4.4)	(6.6)	(2.2)				(100.0)				
総計	33	27	2,139	20,105	40,899	28,772	11,484	3,950	1,455	506	198	129	109,697	2,818	112,515	59.43	12.05
(%)	(0.0)	(0.0)	(1.9)	(18.3)	(37.3)	(26.2)	(10.5)	(3.6)	(1.3)	(0.5)	(0.2)	(0.1)	(100.0)				
平均	6.67	1.64	1.79	2.19	2.51	2.82	3.21	3.64	4.12	4.54	4.89	5.23	2.67		2.67		
標準偏差	10.77	1.15	0.97	1.01	0.95	0.99	1.05	1.12	1.14	1.32	1.37	1.44	1.09		1.09		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査