

表 2464 透析後体重(kg) 体重減少量(kg)別（施設血液透析患者 女性のみ 透析回数週3回 透析歴2年以上）

体重減少量	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
0.0未満		7	105	102	63	16	4			1			298		298	44.70	10.39
(%)		(2.3)	(35.2)	(34.2)	(21.1)	(5.4)	(1.3)			(0.3)			(100.0)				
0.0～		138	1,347	1,679	595	144	39	6	1			1	3,950		3,950	43.43	8.99
(%)		(3.5)	(34.1)	(42.5)	(15.1)	(3.6)	(1.0)	(0.2)	(0.0)			(0.0)	(100.0)				
1.0～		352	6,171	9,459	3,747	865	142	32	10	2			20,780		20,780	44.55	8.52
(%)		(1.7)	(29.7)	(45.5)	(18.0)	(4.2)	(0.7)	(0.2)	(0.0)	(0.0)			(100.0)				
2.0～		121	4,984	13,043	7,748	2,452	601	144	23	7	2	2	29,127		29,127	48.35	9.46
(%)		(0.4)	(17.1)	(44.8)	(26.6)	(8.4)	(2.1)	(0.5)	(0.1)	(0.0)	(0.0)	(0.0)	(100.0)				
3.0～		5	579	2,869	2,707	1,429	570	193	75	9	6	1	8,443		8,443	54.35	11.82
(%)		(0.1)	(6.9)	(34.0)	(32.1)	(16.9)	(6.8)	(2.3)	(0.9)	(0.1)	(0.1)	(0.0)	(100.0)				
4.0～			24	212	304	256	171	100	40	11	5	3	1,126		1,126	62.97	15.31
(%)			(2.1)	(18.8)	(27.0)	(22.7)	(15.2)	(8.9)	(3.6)	(1.0)	(0.4)	(0.3)	(100.0)				
5.0～		1	7	35	25	24	22	21	7	2	3	2	149		149	65.01	19.96
(%)		(0.7)	(4.7)	(23.5)	(16.8)	(16.1)	(14.8)	(14.1)	(4.7)	(1.3)	(2.0)	(1.3)	(100.0)				
6.0～			1	5	3	1	3	2	2	1	1		19		19	66.93	24.08
(%)			(5.3)	(26.3)	(15.8)	(5.3)	(15.8)	(10.5)	(10.5)	(5.3)	(5.3)		(100.0)				
7.0～			2	5	2	1			1	1			12		12	55.14	21.04
(%)			(16.7)	(41.7)	(16.7)	(8.3)			(8.3)	(8.3)			(100.0)				
8.0～	3	3	10	13	9	4	2						44		44	43.85	15.69
(%)	(6.8)	(6.8)	(22.7)	(29.5)	(20.5)	(9.1)	(4.5)						(100.0)				
合計	3	627	13,230	27,422	15,203	5,192	1,554	498	159	34	17	9	63,948		63,948	47.89	10.44
(%)	(0.0)	(1.0)	(20.7)	(42.9)	(23.8)	(8.1)	(2.4)	(0.8)	(0.2)	(0.1)	(0.0)	(0.0)	(100.0)				
記載なし	26	2	3	6	4	2	3	3	1				50	1,696	1,746	32.11	26.41
(%)	(52.0)	(4.0)	(6.0)	(12.0)	(8.0)	(4.0)	(6.0)	(6.0)	(2.0)				(100.0)				
総計	29	629	13,233	27,428	15,207	5,194	1,557	501	160	34	17	9	63,998	1,696	65,694	47.87	10.47
(%)	(0.0)	(1.0)	(20.7)	(42.9)	(23.8)	(8.1)	(2.4)	(0.8)	(0.3)	(0.1)	(0.0)	(0.0)	(100.0)				
平均	21.27	1.46	1.81	2.10	2.34	2.62	2.94	3.29	3.55	3.61	4.02	3.67	2.17		2.17		
標準偏差	8.44	0.96	0.78	0.81	0.83	0.91	0.98	0.98	1.00	1.37	1.00	1.44	0.89		0.89		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査