

表 2471 透析後体重(kg) 透析前BUN(mg/dL)別（施設血液透析患者 透析回数週3回 透析歴2年以上）

透析前BUN	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
20未満	3	7	96	167	146	35	15	2	2				473	6	479	48.58	10.79
(%)	(0.6)	(1.5)	(20.3)	(35.3)	(30.9)	(7.4)	(3.2)	(0.4)	(0.4)				(100.0)				
20～		28	417	927	687	273	95	19	6	1			2,453	27	2,480	49.68	10.94
(%)		(1.1)	(17.0)	(37.8)	(28.0)	(11.1)	(3.9)	(0.8)	(0.2)	(0.0)			(100.0)				
30～	5	69	1,347	3,303	3,260	1,572	507	161	45	14	3	2	10,288	70	10,358	52.08	11.71
(%)	(0.0)	(0.7)	(13.1)	(32.1)	(31.7)	(15.3)	(4.9)	(1.6)	(0.4)	(0.1)	(0.0)	(0.0)	(100.0)				
40～	11	120	2,627	7,928	8,702	4,834	1,683	540	157	52	19	14	26,687	124	26,811	53.93	12.11
(%)	(0.0)	(0.4)	(9.8)	(29.7)	(32.6)	(18.1)	(6.3)	(2.0)	(0.6)	(0.2)	(0.1)	(0.1)	(100.0)				
50～	15	158	3,548	11,549	14,010	8,427	3,255	945	347	123	28	27	42,432	227	42,659	55.15	12.41
(%)	(0.0)	(0.4)	(8.4)	(27.2)	(33.0)	(19.9)	(7.7)	(2.2)	(0.8)	(0.3)	(0.1)	(0.1)	(100.0)				
60～	17	133	3,329	11,050	13,823	8,773	3,433	1,200	449	150	55	29	42,441	202	42,643	55.89	12.83
(%)	(0.0)	(0.3)	(7.8)	(26.0)	(32.6)	(20.7)	(8.1)	(2.8)	(1.1)	(0.4)	(0.1)	(0.1)	(100.0)				
70～	8	71	2,212	7,219	9,022	5,878	2,371	900	346	91	54	26	28,198	124	28,322	56.27	13.14
(%)	(0.0)	(0.3)	(7.8)	(25.6)	(32.0)	(20.8)	(8.4)	(3.2)	(1.2)	(0.3)	(0.2)	(0.1)	(100.0)				
80～		33	1,077	3,470	4,083	2,672	1,103	424	150	66	33	24	13,135	77	13,212	56.28	13.62
(%)		(0.3)	(8.2)	(26.4)	(31.1)	(20.3)	(8.4)	(3.2)	(1.1)	(0.5)	(0.3)	(0.2)	(100.0)				
90～	1	22	433	1,204	1,516	962	352	171	74	27	12	6	4,780	23	4,803	56.18	14.00
(%)	(0.0)	(0.5)	(9.1)	(25.2)	(31.7)	(20.1)	(7.4)	(3.6)	(1.5)	(0.6)	(0.3)	(0.1)	(100.0)				
100～	1	4	120	324	407	268	120	47	23	11	4	8	1,337	11	1,348	57.09	15.06
(%)	(0.1)	(0.3)	(9.0)	(24.2)	(30.4)	(20.0)	(9.0)	(3.5)	(1.7)	(0.8)	(0.3)	(0.6)	(100.0)				
110～		4	34	93	108	88	37	13	9	2	4	1	393	1	394	57.58	15.99
(%)		(1.0)	(8.7)	(23.7)	(27.5)	(22.4)	(9.4)	(3.3)	(2.3)	(0.5)	(1.0)	(0.3)	(100.0)				
120～	1	2	28	50	62	34	14	7	3		2		203	1	204	54.47	14.95
(%)	(0.5)	(1.0)	(13.8)	(24.6)	(30.5)	(16.7)	(6.9)	(3.4)	(1.5)		(1.0)		(100.0)				
合計	62	651	15,268	47,284	55,826	33,816	12,985	4,429	1,611	537	214	137	172,820	893	173,713	55.18	12.78
(%)	(0.0)	(0.4)	(8.8)	(27.4)	(32.3)	(19.6)	(7.5)	(2.6)	(0.9)	(0.3)	(0.1)	(0.1)	(100.0)				
記載なし		5	104	249	280	150	56	22	4	3	1	1	875	3,621	4,496	53.79	12.75
(%)		(0.6)	(11.9)	(28.5)	(32.0)	(17.1)	(6.4)	(2.5)	(0.5)	(0.3)	(0.1)	(0.1)	(100.0)				
総計	62	656	15,372	47,533	56,106	33,966	13,041	4,451	1,615	540	215	138	173,695	4,514	178,209	55.17	12.78
(%)	(0.0)	(0.4)	(8.8)	(27.4)	(32.3)	(19.6)	(7.5)	(2.6)	(0.9)	(0.3)	(0.1)	(0.1)	(100.0)				
平均	57.19	56.93	58.90	59.84	60.86	62.07	62.82	64.51	65.63	66.04	70.11	69.56	60.95	59.76	60.94		
標準偏差	18.82	18.06	16.98	15.96	15.39	14.92	14.81	15.11	15.25	15.43	16.22	16.72	15.63	16.74	15.64		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査