

表 2483 透析後体重(kg) %クレアチニン産生速度(%)別（施設血液透析患者 男性のみ 透析回数週3回 透析歴2年以上）

%クレアチニン 産生速度	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
60未満		8	638	2,505	2,361	1,217	447	149	60	18	12	9	7,424		7,424	53.88	12.29
(%)		(0.1)	(8.6)	(33.7)	(31.8)	(16.4)	(6.0)	(2.0)	(0.8)	(0.2)	(0.2)	(0.1)	(100.0)				
60～		5	230	1,628	2,120	1,098	428	159	79	21	10	8	5,786		5,786	56.47	12.38
(%)		(0.1)	(4.0)	(28.1)	(36.6)	(19.0)	(7.4)	(2.7)	(1.4)	(0.4)	(0.2)	(0.1)	(100.0)				
70～		2	276	2,191	3,261	1,945	791	287	94	37	13	12	8,909		8,909	57.59	12.26
(%)		(0.0)	(3.1)	(24.6)	(36.6)	(21.8)	(8.9)	(3.2)	(1.1)	(0.4)	(0.1)	(0.1)	(100.0)				
80～		3	267	2,646	4,783	2,942	1,160	449	196	68	26	19	12,559		12,559	58.76	12.44
(%)		(0.0)	(2.1)	(21.1)	(38.1)	(23.4)	(9.2)	(3.6)	(1.6)	(0.5)	(0.2)	(0.2)	(100.0)				
90～		1	211	2,990	6,069	4,150	1,599	582	233	91	33	30	15,989		15,989	59.61	12.16
(%)		(0.0)	(1.3)	(18.7)	(38.0)	(26.0)	(10.0)	(3.6)	(1.5)	(0.6)	(0.2)	(0.2)	(100.0)				
100～			161	2,666	6,496	4,593	1,838	661	254	99	30	20	16,818		16,818	60.36	11.88
(%)			(1.0)	(15.9)	(38.6)	(27.3)	(10.9)	(3.9)	(1.5)	(0.6)	(0.2)	(0.1)	(100.0)				
110～			93	1,969	5,769	4,417	1,836	647	217	71	28	16	15,063		15,063	61.15	11.53
(%)			(0.6)	(13.1)	(38.3)	(29.3)	(12.2)	(4.3)	(1.4)	(0.5)	(0.2)	(0.1)	(100.0)				
120～	1		53	1,250	4,017	3,449	1,419	419	144	38	19	10	10,819		10,819	61.50	11.08
(%)	(0.0)		(0.5)	(11.6)	(37.1)	(31.9)	(13.1)	(3.9)	(1.3)	(0.4)	(0.2)	(0.1)	(100.0)				
130～			23	624	2,251	2,040	816	237	78	23	6	1	6,099		6,099	61.81	10.62
(%)			(0.4)	(10.2)	(36.9)	(33.4)	(13.4)	(3.9)	(1.3)	(0.4)	(0.1)	(0.0)	(100.0)				
140～	2		11	370	1,501	1,513	541	163	32	10	7	1	4,151		4,151	61.84	10.28
(%)	(0.0)		(0.3)	(8.9)	(36.2)	(36.4)	(13.0)	(3.9)	(0.8)	(0.2)	(0.2)	(0.0)	(100.0)				
合計	3	19	1,963	18,839	38,628	27,364	10,875	3,753	1,387	476	184	126	103,617		103,617	59.51	12.02
(%)	(0.0)	(0.0)	(1.9)	(18.2)	(37.3)	(26.4)	(10.5)	(3.6)	(1.3)	(0.5)	(0.2)	(0.1)	(100.0)				
記載なし	30	8	176	1,266	2,271	1,408	609	197	68	30	14	3	6,080	2,818	8,898	58.13	12.58
(%)	(0.5)	(0.1)	(2.9)	(20.8)	(37.4)	(23.2)	(10.0)	(3.2)	(1.1)	(0.5)	(0.2)	(0.0)	(100.0)				
総計	33	27	2,139	20,105	40,899	28,772	11,484	3,950	1,455	506	198	129	109,697	2,818	112,515	59.43	12.05
(%)	(0.0)	(0.0)	(1.9)	(18.3)	(37.3)	(26.2)	(10.5)	(3.6)	(1.3)	(0.5)	(0.2)	(0.1)	(100.0)				
平均	204.77	62.27	73.24	89.81	99.67	103.76	104.00	102.44	99.77	99.15	99.01	92.90	99.00		99.00		
標準偏差	75.87	19.12	28.30	26.60	24.81	24.34	23.77	23.44	22.43	20.93	28.00	21.90	25.64		25.64		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査