

表 2484 透析後体重(kg) %クレアチニン産生速度(%)別（施設血液透析患者 女性のみ 透析回数週3回 透析歴2年以上）

%クレアチニン 産生速度	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
60未満		252	2,051	1,781	694	230	76	25	7	5	3	2	5,126		5,126	43.02	10.64
(%)		(4.9)	(40.0)	(34.7)	(13.5)	(4.5)	(1.5)	(0.5)	(0.1)	(0.1)	(0.1)	(0.0)	(100.0)				
60～		72	1,026	1,332	571	245	58	25	6	2	2		3,339		3,339	45.59	10.78
(%)		(2.2)	(30.7)	(39.9)	(17.1)	(7.3)	(1.7)	(0.7)	(0.2)	(0.1)	(0.1)		(100.0)				
70～		58	1,339	2,090	1,006	353	126	50	15	4	2	2	5,045		5,045	46.88	11.02
(%)		(1.1)	(26.5)	(41.4)	(19.9)	(7.0)	(2.5)	(1.0)	(0.3)	(0.1)	(0.0)	(0.0)	(100.0)				
80～		65	1,623	2,965	1,566	606	197	63	28	6	4	1	7,124		7,124	47.86	11.04
(%)		(0.9)	(22.8)	(41.6)	(22.0)	(8.5)	(2.8)	(0.9)	(0.4)	(0.1)	(0.1)	(0.0)	(100.0)				
90～		45	1,780	3,836	2,098	770	230	80	27	2	1		8,869		8,869	48.20	10.40
(%)		(0.5)	(20.1)	(43.3)	(23.7)	(8.7)	(2.6)	(0.9)	(0.3)	(0.0)	(0.0)		(100.0)				
100～		34	1,680	4,335	2,484	848	274	82	30	6	4		9,777		9,777	48.79	10.27
(%)		(0.3)	(17.2)	(44.3)	(25.4)	(8.7)	(2.8)	(0.8)	(0.3)	(0.1)	(0.0)		(100.0)				
110～		23	1,388	3,925	2,418	742	219	56	14	4		2	8,791		8,791	48.88	9.69
(%)		(0.3)	(15.8)	(44.6)	(27.5)	(8.4)	(2.5)	(0.6)	(0.2)	(0.0)		(0.0)	(100.0)				
120～		12	861	3,040	1,813	581	148	47	14	3			6,519		6,519	49.30	9.52
(%)		(0.2)	(13.2)	(46.6)	(27.8)	(8.9)	(2.3)	(0.7)	(0.2)	(0.0)			(100.0)				
130～		5	412	1,592	1,065	324	83	22	4		1	1	3,509		3,509	49.60	9.21
(%)		(0.1)	(11.7)	(45.4)	(30.4)	(9.2)	(2.4)	(0.6)	(0.1)		(0.0)	(0.0)	(100.0)				
140～	2	7	232	1,024	701	240	70	12	7	2		1	2,298		2,298	50.30	9.85
(%)	(0.1)	(0.3)	(10.1)	(44.6)	(30.5)	(10.4)	(3.0)	(0.5)	(0.3)	(0.1)		(0.0)	(100.0)				
合計	2	573	12,392	25,920	14,416	4,939	1,481	462	152	34	17	9	60,397		60,397	47.94	10.44
(%)	(0.0)	(0.9)	(20.5)	(42.9)	(23.9)	(8.2)	(2.5)	(0.8)	(0.3)	(0.1)	(0.0)	(0.0)	(100.0)				
記載なし	27	56	841	1,508	791	255	76	39	8				3,601	1,696	5,297	46.77	11.03
(%)	(0.7)	(1.6)	(23.4)	(41.9)	(22.0)	(7.1)	(2.1)	(1.1)	(0.2)				(100.0)				
総計	29	629	13,233	27,428	15,207	5,194	1,557	501	160	34	17	9	63,998	1,696	65,694	47.87	10.47
(%)	(0.0)	(1.0)	(20.7)	(42.9)	(23.8)	(8.1)	(2.4)	(0.8)	(0.3)	(0.1)	(0.0)	(0.0)	(100.0)				
平均	165.00	67.10	88.40	100.02	103.42	102.10	101.07	97.83	97.80	92.69	80.32	94.34	98.30		98.30		
標準偏差	8.06	30.06	28.14	25.57	24.28	24.56	24.60	24.00	22.99	28.37	30.39	35.13	26.49		26.49		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査