

表 2485 透析後体重(kg) 透析前アルブミン濃度(g/dL)別（施設血液透析患者 透析回数週3回）

透析前アルブミン濃度	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
1.5未満		1	23	43	41	21	7	3	1	1			141	5	146	52.17	13.17
(%)		(0.7)	(16.3)	(30.5)	(29.1)	(14.9)	(5.0)	(2.1)	(0.7)	(0.7)			(100.0)				
1.5～	1	7	123	281	221	81	27	6	4	4	1		756	15	771	50.04	11.92
(%)	(0.1)	(0.9)	(16.3)	(37.2)	(29.2)	(10.7)	(3.6)	(0.8)	(0.5)	(0.5)	(0.1)		(100.0)				
2.0～	3	40	675	1,275	746	270	71	29	8	3	3		3,123	56	3,179	47.80	10.94
(%)	(0.1)	(1.3)	(21.6)	(40.8)	(23.9)	(8.6)	(2.3)	(0.9)	(0.3)	(0.1)	(0.1)		(100.0)				
2.5～	8	152	2,298	4,504	3,360	1,186	355	99	28	13	6	4	12,013	137	12,150	48.98	11.08
(%)	(0.1)	(1.3)	(19.1)	(37.5)	(28.0)	(9.9)	(3.0)	(0.8)	(0.2)	(0.1)	(0.0)	(0.0)	(100.0)				
3.0～	21	284	5,976	16,360	16,948	8,314	2,602	829	267	88	37	28	51,754	328	52,082	52.75	11.83
(%)	(0.0)	(0.5)	(11.5)	(31.6)	(32.7)	(16.1)	(5.0)	(1.6)	(0.5)	(0.2)	(0.1)	(0.1)	(100.0)				
3.5～	34	227	7,325	26,874	35,691	22,731	8,788	3,096	1,108	403	163	111	106,551	584	107,135	56.38	12.79
(%)	(0.0)	(0.2)	(6.9)	(25.2)	(33.5)	(21.3)	(8.2)	(2.9)	(1.0)	(0.4)	(0.2)	(0.1)	(100.0)				
4.0～	15	59	2,043	8,461	12,553	9,768	4,606	1,758	665	211	80	37	40,256	250	40,506	58.87	13.46
(%)	(0.0)	(0.1)	(5.1)	(21.0)	(31.2)	(24.3)	(11.4)	(4.4)	(1.7)	(0.5)	(0.2)	(0.1)	(100.0)				
4.5～	1	4	112	438	735	585	287	115	52	14	2	4	2,349	13	2,362	59.80	13.82
(%)	(0.0)	(0.2)	(4.8)	(18.6)	(31.3)	(24.9)	(12.2)	(4.9)	(2.2)	(0.6)	(0.1)	(0.2)	(100.0)				
5.0～		1	12	51	52	26	18	11	1				172		172	56.11	13.23
(%)		(0.6)	(7.0)	(29.7)	(30.2)	(15.1)	(10.5)	(6.4)	(0.6)				(100.0)				
合計	83	775	18,587	58,287	70,347	42,982	16,761	5,946	2,134	737	292	184	217,115	1,388	218,503	55.46	12.90
(%)	(0.0)	(0.4)	(8.6)	(26.8)	(32.4)	(19.8)	(7.7)	(2.7)	(1.0)	(0.3)	(0.1)	(0.1)	(100.0)				
記載なし	1	27	530	1,522	1,762	1,100	423	135	61	18	7	5	5,591	4,877	10,468	55.05	12.89
(%)	(0.0)	(0.5)	(9.5)	(27.2)	(31.5)	(19.7)	(7.6)	(2.4)	(1.1)	(0.3)	(0.1)	(0.1)	(100.0)				
総計	84	802	19,117	59,809	72,109	44,082	17,184	6,081	2,195	755	299	189	222,706	6,265	228,971	55.45	12.90
(%)	(0.0)	(0.4)	(8.6)	(26.9)	(32.4)	(19.8)	(7.7)	(2.7)	(1.0)	(0.3)	(0.1)	(0.1)	(100.0)				
平均	3.49	3.26	3.41	3.52	3.61	3.69	3.75	3.77	3.80	3.76	3.74	3.72	3.60	3.49	3.60		
標準偏差	0.54	0.52	0.50	0.46	0.42	0.39	0.38	0.38	0.37	0.40	0.37	0.32	0.44	0.55	0.44		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査