

表 2488 透析後体重(kg) 透析前リン濃度(mg/dL)別(施設血液透析患者 透析回数週3回)

透析前リン濃度	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
3未満	6	104	1,615	3,190	2,479	1,036	309	109	26	7	3	2	8,886	75	8,961	49.75	11.62
(%)	(0.1)	(1.2)	(18.2)	(35.9)	(27.9)	(11.7)	(3.5)	(1.2)	(0.3)	(0.1)	(0.0)	(0.0)	(100.0)				
3.0～	4	84	1,396	3,480	3,585	1,653	511	121	46	14	4	3	10,901	79	10,980	52.00	11.41
(%)	(0.0)	(0.8)	(12.8)	(31.9)	(32.9)	(15.2)	(4.7)	(1.1)	(0.4)	(0.1)	(0.0)	(0.0)	(100.0)				
3.5～	9	90	2,065	5,795	6,162	3,257	1,105	278	81	35	9	5	18,891	127	19,018	53.15	11.70
(%)	(0.0)	(0.5)	(10.9)	(30.7)	(32.6)	(17.2)	(5.8)	(1.5)	(0.4)	(0.2)	(0.0)	(0.0)	(100.0)				
4.0～	13	107	2,653	8,273	9,344	5,155	1,699	551	158	42	24	8	28,027	165	28,192	53.95	11.85
(%)	(0.0)	(0.4)	(9.5)	(29.5)	(33.3)	(18.4)	(6.1)	(2.0)	(0.6)	(0.1)	(0.1)	(0.0)	(100.0)				
4.5～	12	110	2,806	9,190	11,163	6,455	2,414	754	247	60	27	13	33,251	217	33,468	54.91	12.16
(%)	(0.0)	(0.3)	(8.4)	(27.6)	(33.6)	(19.4)	(7.3)	(2.3)	(0.7)	(0.2)	(0.1)	(0.0)	(100.0)				
5.0～	16	100	2,606	8,936	11,124	6,961	2,579	858	273	95	31	15	33,594	215	33,809	55.53	12.41
(%)	(0.0)	(0.3)	(7.8)	(26.6)	(33.1)	(20.7)	(7.7)	(2.6)	(0.8)	(0.3)	(0.1)	(0.0)	(100.0)				
5.5～	13	64	2,045	7,411	9,420	6,162	2,391	839	328	117	38	28	28,856	156	29,012	56.33	12.93
(%)	(0.0)	(0.2)	(7.1)	(25.7)	(32.6)	(21.4)	(8.3)	(2.9)	(1.1)	(0.4)	(0.1)	(0.1)	(100.0)				
6.0～		6	349	1,166	1,665	1,070	402	157	53	23	3	3	4,897	31	4,928	56.63	12.70
(%)		(0.1)	(7.1)	(23.8)	(34.0)	(21.9)	(8.2)	(3.2)	(1.1)	(0.5)	(0.1)	(0.1)	(100.0)				
6.1～	4	37	1,094	3,930	5,273	3,583	1,550	619	214	68	29	17	16,418	87	16,505	57.25	13.27
(%)	(0.0)	(0.2)	(6.7)	(23.9)	(32.1)	(21.8)	(9.4)	(3.8)	(1.3)	(0.4)	(0.2)	(0.1)	(100.0)				
6.5～	3	27	905	3,195	4,610	3,230	1,454	536	217	77	33	17	14,304	73	14,377	57.87	13.57
(%)	(0.0)	(0.2)	(6.3)	(22.3)	(32.2)	(22.6)	(10.2)	(3.7)	(1.5)	(0.5)	(0.2)	(0.1)	(100.0)				
7.0～	2	17	524	1,900	2,787	1,931	989	392	159	59	19	22	8,801	59	8,860	58.65	14.18
(%)	(0.0)	(0.2)	(6.0)	(21.6)	(31.7)	(21.9)	(11.2)	(4.5)	(1.8)	(0.7)	(0.2)	(0.2)	(100.0)				
7.5～		14	343	1,121	1,530	1,235	606	286	122	40	11	7	5,315	44	5,359	59.18	14.54
(%)		(0.3)	(6.5)	(21.1)	(28.8)	(23.2)	(11.4)	(5.4)	(2.3)	(0.8)	(0.2)	(0.1)	(100.0)				
8.0～		10	167	631	861	764	398	201	72	31	23	17	3,175	17	3,192	60.79	15.64
(%)		(0.3)	(5.3)	(19.9)	(27.1)	(24.1)	(12.5)	(6.3)	(2.3)	(1.0)	(0.7)	(0.5)	(100.0)				
8.5～	1	5	112	392	557	475	245	113	56	32	13	12	2,013	10	2,023	61.04	16.26
(%)	(0.0)	(0.2)	(5.6)	(19.5)	(27.7)	(23.6)	(12.2)	(5.6)	(2.8)	(1.6)	(0.6)	(0.6)	(100.0)				
9.0～	1	10	196	540	846	723	381	214	122	48	29	19	3,129	27	3,156	61.88	16.97
(%)	(0.0)	(0.3)	(6.3)	(17.3)	(27.0)	(23.1)	(12.2)	(6.8)	(3.9)	(1.5)	(0.9)	(0.6)	(100.0)				
合計	84	785	18,876	59,150	71,406	43,690	17,033	6,028	2,174	748	296	188	220,458	1,382	221,840	55.46	12.90
(%)	(0.0)	(0.4)	(8.6)	(26.8)	(32.4)	(19.8)	(7.7)	(2.7)	(1.0)	(0.3)	(0.1)	(0.1)	(100.0)				
記載なし		17	241	659	703	392	151	53	21	7	3	1	2,248	4,883	7,131	53.97	12.77
(%)		(0.8)	(10.7)	(29.3)	(31.3)	(17.4)	(6.7)	(2.4)	(0.9)	(0.3)	(0.1)	(0.0)	(100.0)				
総計	84	802	19,117	59,809	72,109	44,082	17,184	6,081	2,195	755	299	189	222,706	6,265	228,971	55.45	12.90
(%)	(0.0)	(0.4)	(8.6)	(26.9)	(32.4)	(19.8)	(7.7)	(2.7)	(1.0)	(0.3)	(0.1)	(0.1)	(100.0)				
平均	4.83	4.57	4.82	5.00	5.18	5.37	5.57	5.81	6.04	6.20	6.40	6.63	5.20	5.13	5.20		
標準偏差	1.35	1.58	1.47	1.38	1.37	1.40	1.45	1.55	1.63	1.66	1.94	1.82	1.43	1.54	1.43		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査