

表 2490 透析後体重(kg) PTH値(pg/mL)別(施設血液透析患者 透析回数週3回) PTH測定法:whole-PTH

PTH値	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
20未満	1	6	193	463	435	268	87	28	8	7	1		1,497	2	1,499	53.10	12.44
(%)	(0.1)	(0.4)	(12.9)	(30.9)	(29.1)	(17.9)	(5.8)	(1.9)	(0.5)	(0.5)	(0.1)		(100.0)				
20～		12	194	539	605	316	121	39	13	2		2	1,843	5	1,848	53.83	12.44
(%)		(0.7)	(10.5)	(29.2)	(32.8)	(17.1)	(6.6)	(2.1)	(0.7)	(0.1)		(0.1)	(100.0)				
35～	2	9	161	443	531	288	113	47	9	3		1	1,607	6	1,613	54.40	12.41
(%)	(0.1)	(0.6)	(10.0)	(27.6)	(33.0)	(17.9)	(7.0)	(2.9)	(0.6)	(0.2)		(0.1)	(100.0)				
50～	2	5	186	639	794	517	211	74	6	8	5	3	2,450	9	2,459	55.86	12.62
(%)	(0.1)	(0.2)	(7.6)	(26.1)	(32.4)	(21.1)	(8.6)	(3.0)	(0.2)	(0.3)	(0.2)	(0.1)	(100.0)				
75～	1	3	174	489	744	456	160	72	25	6	5	4	2,139	9	2,148	56.47	13.21
(%)	(0.0)	(0.1)	(8.1)	(22.9)	(34.8)	(21.3)	(7.5)	(3.4)	(1.2)	(0.3)	(0.2)	(0.2)	(100.0)				
100～		3	115	438	577	391	168	54	21	6	1	2	1,776	11	1,787	56.84	12.76
(%)		(0.2)	(6.5)	(24.7)	(32.5)	(22.0)	(9.5)	(3.0)	(1.2)	(0.3)	(0.1)	(0.1)	(100.0)				
125～		5	70	257	404	272	118	30	19	3	1		1,179	8	1,187	57.33	12.64
(%)		(0.4)	(5.9)	(21.8)	(34.3)	(23.1)	(10.0)	(2.5)	(1.6)	(0.3)	(0.1)		(100.0)				
150～			1	8	13	9	4	1	1				37		37	58.19	12.36
(%)			(2.7)	(21.6)	(35.1)	(24.3)	(10.8)	(2.7)	(2.7)				(100.0)				
151～		3	70	205	321	197	94	28	7	4	1	1	931	3	934	56.89	12.95
(%)		(0.3)	(7.5)	(22.0)	(34.5)	(21.2)	(10.1)	(3.0)	(0.8)	(0.4)	(0.1)	(0.1)	(100.0)				
180～		2	46	168	194	148	66	24	12	1	2		663	3	666	57.30	13.57
(%)		(0.3)	(6.9)	(25.3)	(29.3)	(22.3)	(10.0)	(3.6)	(1.8)	(0.2)	(0.3)		(100.0)				
210～		2	29	109	132	112	53	17	13	2	1		470	2	472	58.48	13.75
(%)		(0.4)	(6.2)	(23.2)	(28.1)	(23.8)	(11.3)	(3.6)	(2.8)	(0.4)	(0.2)		(100.0)				
240～		1	31	139	160	116	44	20	9	2	1	2	525	1	526	57.13	13.91
(%)		(0.2)	(5.9)	(26.5)	(30.5)	(22.1)	(8.4)	(3.8)	(1.7)	(0.4)	(0.2)	(0.4)	(100.0)				
300～			14	39	71	39	17	15	7	1	2		205		205	59.68	15.15
(%)			(6.8)	(19.0)	(34.6)	(19.0)	(8.3)	(7.3)	(3.4)	(0.5)	(1.0)		(100.0)				
350～		2	9	31	45	20	15	8	3	1			134	1	135	57.33	14.68
(%)		(1.5)	(6.7)	(23.1)	(33.6)	(14.9)	(11.2)	(6.0)	(2.2)	(0.7)			(100.0)				
400～			18	38	46	28	13	6	1		1	1	152		152	55.98	15.23
(%)			(11.8)	(25.0)	(30.3)	(18.4)	(8.6)	(3.9)	(0.7)		(0.7)	(0.7)	(100.0)				
500～		1	18	47	49	31	12	6	1	1			166		166	54.95	12.90
(%)		(0.6)	(10.8)	(28.3)	(29.5)	(18.7)	(7.2)	(3.6)	(0.6)	(0.6)			(100.0)				
合計	6	54	1,329	4,052	5,121	3,208	1,296	469	155	47	21	16	15,774	60	15,834	55.82	12.97
(%)	(0.0)	(0.3)	(8.4)	(25.7)	(32.5)	(20.3)	(8.2)	(3.0)	(1.0)	(0.3)	(0.1)	(0.1)	(100.0)				
記載なし		5	59	135	116	69	13	5	3				405	33	438	51.51	11.39
(%)		(1.2)	(14.6)	(33.3)	(28.6)	(17.0)	(3.2)	(1.2)	(0.7)				(100.0)				
総計	6	59	1,388	4,187	5,237	3,277	1,309	474	158	47	21	16	16,179	93	16,272	55.71	12.95
(%)	(0.0)	(0.4)	(8.6)	(25.9)	(32.4)	(20.3)	(8.1)	(2.9)	(1.0)	(0.3)	(0.1)	(0.1)	(100.0)				
平均	52.00	104.65	97.82	101.66	104.31	107.84	115.03	122.40	142.82	117.74	151.00	129.44	105.70	104.02	105.70		
標準偏差	21.71	140.55	131.11	120.44	101.41	93.93	107.46	116.73	110.66	122.48	111.98	124.80	109.37	65.83	109.24		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査