

表 2500 透析後体重(kg) 透析歴(年)別 (血液透析濾過患者 透析回数週3回)

透析歴	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
2年未満	1	9	286	1,048	1,479	1,132	512	199	84	33	15	6	4,804	160	4,964	58.59	13.88
(%)	(0.0)	(0.2)	(6.0)	(21.8)	(30.8)	(23.6)	(10.7)	(4.1)	(1.7)	(0.7)	(0.3)	(0.1)	(100.0)				
2年～	4	22	420	1,745	2,600	2,013	908	417	180	51	28	16	8,404	211	8,615	59.38	14.27
(%)	(0.0)	(0.3)	(5.0)	(20.8)	(30.9)	(24.0)	(10.8)	(5.0)	(2.1)	(0.6)	(0.3)	(0.2)	(100.0)				
5年～	1	12	588	2,274	3,099	2,300	1,003	414	118	43	8	12	9,872	244	10,116	57.84	13.17
(%)	(0.0)	(0.1)	(6.0)	(23.0)	(31.4)	(23.3)	(10.2)	(4.2)	(1.2)	(0.4)	(0.1)	(0.1)	(100.0)				
10年～	1	10	421	1,549	1,965	1,382	530	134	51	10	7	2	6,062	132	6,194	56.09	12.11
(%)	(0.0)	(0.2)	(6.9)	(25.6)	(32.4)	(22.8)	(8.7)	(2.2)	(0.8)	(0.2)	(0.1)	(0.0)	(100.0)				
15年～	3	8	298	1,196	1,317	722	235	65	14	4	3		3,865	59	3,924	54.07	11.17
(%)	(0.1)	(0.2)	(7.7)	(30.9)	(34.1)	(18.7)	(6.1)	(1.7)	(0.4)	(0.1)	(0.1)		(100.0)				
20年～	1	4	259	903	869	404	114	22	6			1	2,583	54	2,637	52.22	10.47
(%)	(0.0)	(0.2)	(10.0)	(35.0)	(33.6)	(15.6)	(4.4)	(0.9)	(0.2)			(0.0)	(100.0)				
25年～	1	5	168	606	562	204	66	13	1	1			1,627	24	1,651	51.30	10.08
(%)	(0.1)	(0.3)	(10.3)	(37.2)	(34.5)	(12.5)	(4.1)	(0.8)	(0.1)	(0.1)			(100.0)				
30年～		5	135	405	373	135	37	4	1				1,095	28	1,123	50.61	9.64
(%)		(0.5)	(12.3)	(37.0)	(34.1)	(12.3)	(3.4)	(0.4)	(0.1)				(100.0)				
35年～		1	80	227	192	57	15	4					576	11	587	49.88	9.38
(%)		(0.2)	(13.9)	(39.4)	(33.3)	(9.9)	(2.6)	(0.7)					(100.0)				
40年～			27	48	31	14	2						122	2	124	48.58	9.18
(%)			(22.1)	(39.3)	(25.4)	(11.5)	(1.6)						(100.0)				
合計	12	76	2,682	10,001	12,487	8,363	3,422	1,272	455	142	61	37	39,010	925	39,935	56.63	13.03
(%)	(0.0)	(0.2)	(6.9)	(25.6)	(32.0)	(21.4)	(8.8)	(3.3)	(1.2)	(0.4)	(0.2)	(0.1)	(100.0)				
不明				4	3	2	1	1					11		11	58.34	13.29
(%)				(36.4)	(27.3)	(18.2)	(9.1)	(9.1)					(100.0)				
記載なし																	
(%)																	
総計	12	76	2,682	10,005	12,490	8,365	3,423	1,273	455	142	61	37	39,021	925	39,946	56.63	13.03
(%)	(0.0)	(0.2)	(6.9)	(25.6)	(32.0)	(21.4)	(8.8)	(3.3)	(1.2)	(0.4)	(0.2)	(0.1)	(100.0)				
平均	10.92	10.50	12.62	11.89	10.35	8.59	7.58	6.17	5.24	4.78	4.56	4.84	10.05	8.82	10.02		
標準偏差	9.31	10.01	10.27	9.72	8.96	7.65	6.85	5.62	4.71	4.11	4.32	4.00	8.87	8.60	8.87		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査