

表 2502 透析後体重(kg) 医療機関別（血液透析濾過患者 透析回数週3回）

医療機関	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
国公立大学			9	6	19	7		1					42		42	51.70	10.99
(%)			(21.4)	(14.3)	(45.2)	(16.7)		(2.4)					(100.0)				
私立大学		1	12	64	46	24	14	6	2		1	1	171	2	173	54.74	14.58
(%)		(0.6)	(7.0)	(37.4)	(26.9)	(14.0)	(8.2)	(3.5)	(1.2)		(0.6)	(0.6)	(100.0)				
国立		1	5	11	10	4	3	1	1				36		36	52.59	14.45
(%)		(2.8)	(13.9)	(30.6)	(27.8)	(11.1)	(8.3)	(2.8)	(2.8)				(100.0)				
県市町村立	1	4	113	475	578	332	137	51	18	3	1		1,713	60	1,773	55.79	12.30
(%)	(0.1)	(0.2)	(6.6)	(27.7)	(33.7)	(19.4)	(8.0)	(3.0)	(1.1)	(0.2)	(0.1)		(100.0)				
社会保険	1		25	110	93	50	20	7	3	1	1		311	8	319	54.05	12.45
(%)	(0.3)		(8.0)	(35.4)	(29.9)	(16.1)	(6.4)	(2.3)	(1.0)	(0.3)	(0.3)		(100.0)				
厚生連	1	1	77	255	322	204	66	24	12	7	1	1	971	42	1,013	55.77	13.15
(%)	(0.1)	(0.1)	(7.9)	(26.3)	(33.2)	(21.0)	(6.8)	(2.5)	(1.2)	(0.7)	(0.1)	(0.1)	(100.0)				
その他公的	1	2	62	188	186	132	49	35	4	2	1	1	663	31	694	55.73	13.91
(%)	(0.2)	(0.3)	(9.4)	(28.4)	(28.1)	(19.9)	(7.4)	(5.3)	(0.6)	(0.3)	(0.2)	(0.2)	(100.0)				
私立総合	1	3	69	271	360	245	82	29	9	4	3		1,076	17	1,093	56.20	12.55
(%)	(0.1)	(0.3)	(6.4)	(25.2)	(33.5)	(22.8)	(7.6)	(2.7)	(0.8)	(0.4)	(0.3)		(100.0)				
私立	4	24	779	2,571	3,117	2,009	733	264	91	34	12	8	9,646	369	10,015	55.73	12.73
(%)	(0.0)	(0.2)	(8.1)	(26.7)	(32.3)	(20.8)	(7.6)	(2.7)	(0.9)	(0.4)	(0.1)	(0.1)	(100.0)				
私立診療所	3	40	1,531	6,054	7,759	5,358	2,319	855	315	91	41	26	24,392	396	24,788	57.18	13.15
(%)	(0.0)	(0.2)	(6.3)	(24.8)	(31.8)	(22.0)	(9.5)	(3.5)	(1.3)	(0.4)	(0.2)	(0.1)	(100.0)				
総計	12	76	2,682	10,005	12,490	8,365	3,423	1,273	455	142	61	37	39,021	925	39,946	56.63	13.03
(%)	(0.0)	(0.2)	(6.9)	(25.6)	(32.0)	(21.4)	(8.8)	(3.3)	(1.2)	(0.4)	(0.2)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査