

表 2514 透析後体重(kg) 腎移植の回数別（血液透析濾過患者 透析回数週3回）

腎移植の回数	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
なし	8	58	2,078	7,875	9,790	6,628	2,723	1,029	359	108	49	29	30,734	345	31,079	56.72	13.03
(%)	(0.0)	(0.2)	(6.8)	(25.6)	(31.9)	(21.6)	(8.9)	(3.3)	(1.2)	(0.4)	(0.2)	(0.1)	(100.0)				
1回			48	189	230	122	47	10	3	1			650	6	656	54.46	11.07
(%)			(7.4)	(29.1)	(35.4)	(18.8)	(7.2)	(1.5)	(0.5)	(0.2)			(100.0)				
2回			2	11	13	5	2	1					34	2	36	54.03	10.84
(%)			(5.9)	(32.4)	(38.2)	(14.7)	(5.9)	(2.9)					(100.0)				
3回				1	1								2		2	48.65	10.68
(%)				(50.0)	(50.0)								(100.0)				
4回以上																	
(%)																	
合計	8	58	2,128	8,076	10,034	6,755	2,772	1,040	362	109	49	29	31,420	353	31,773	56.67	13.00
(%)	(0.0)	(0.2)	(6.8)	(25.7)	(31.9)	(21.5)	(8.8)	(3.3)	(1.2)	(0.3)	(0.2)	(0.1)	(100.0)				
不明	1		14	32	47	22	10	6	1	1			134	12	146	55.46	14.21
(%)	(0.7)		(10.4)	(23.9)	(35.1)	(16.4)	(7.5)	(4.5)	(0.7)	(0.7)			(100.0)				
記載なし	3	18	540	1,897	2,409	1,588	641	227	92	32	12	8	7,467	560	8,027	56.48	13.15
(%)	(0.0)	(0.2)	(7.2)	(25.4)	(32.3)	(21.3)	(8.6)	(3.0)	(1.2)	(0.4)	(0.2)	(0.1)	(100.0)				
総計	12	76	2,682	10,005	12,490	8,365	3,423	1,273	455	142	61	37	39,021	925	39,946	56.63	13.03
(%)	(0.0)	(0.2)	(6.9)	(25.6)	(32.0)	(21.4)	(8.8)	(3.3)	(1.2)	(0.4)	(0.2)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査