

表 2520 透析後体重(kg) 体重減少量(kg)別（血液透析濾過患者 透析回数週3回 透析歴2年以上）

体重減少量	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
0.0未満		1	13	26	32	16	7	2					97		97	52.95	12.06
(%)		(1.0)	(13.4)	(26.8)	(33.0)	(16.5)	(7.2)	(2.1)					(100.0)				
0.0～		13	208	480	370	181	52	12	3	1			1,320		1,320	50.55	11.10
(%)		(1.0)	(15.8)	(36.4)	(28.0)	(13.7)	(3.9)	(0.9)	(0.2)	(0.1)			(100.0)				
1.0～		36	1,077	2,813	2,239	963	214	48	9			1	7,400		7,400	50.25	10.20
(%)		(0.5)	(14.6)	(38.0)	(30.3)	(13.0)	(2.9)	(0.6)	(0.1)			(0.0)	(100.0)				
2.0～		10	949	4,436	5,302	2,862	845	198	41	7	3	1	14,654		14,654	54.27	10.52
(%)		(0.1)	(6.5)	(30.3)	(36.2)	(19.5)	(5.8)	(1.4)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
3.0～			132	1,094	2,575	2,473	1,167	405	128	31	5	4	8,014		8,014	61.62	11.95
(%)			(1.6)	(13.7)	(32.1)	(30.9)	(14.6)	(5.1)	(1.6)	(0.4)	(0.1)	(0.0)	(100.0)				
4.0～			12	84	429	635	526	324	117	38	16	10	2,191		2,191	70.33	14.15
(%)			(0.5)	(3.8)	(19.6)	(29.0)	(24.0)	(14.8)	(5.3)	(1.7)	(0.7)	(0.5)	(100.0)				
5.0～			1	5	35	89	92	69	58	28	16	7	400		400	79.74	17.27
(%)			(0.3)	(1.3)	(8.8)	(22.3)	(23.0)	(17.3)	(14.5)	(7.0)	(4.0)	(1.8)	(100.0)				
6.0～					11	7	6	11	13	3	4	4	59		59	84.94	23.71
(%)					(18.6)	(11.9)	(10.2)	(18.6)	(22.0)	(5.1)	(6.8)	(6.8)	(100.0)				
7.0～				3	3	2	1	1	1	1	2	2	16		16	80.15	31.68
(%)				(18.8)	(18.8)	(12.5)	(6.3)	(6.3)	(6.3)	(6.3)	(12.5)	(12.5)	(100.0)				
8.0～		1	1	11	11	2		1	1			2	30		30	57.13	24.35
(%)		(3.3)	(3.3)	(36.7)	(36.7)	(6.7)		(3.3)	(3.3)			(6.7)	(100.0)				
合計		61	2,393	8,952	11,007	7,230	2,910	1,071	371	109	46	31	34,181		34,181	56.37	12.85
(%)		(0.2)	(7.0)	(26.2)	(32.2)	(21.2)	(8.5)	(3.1)	(1.1)	(0.3)	(0.1)	(0.1)	(100.0)				
記載なし	11	6	3	1	1	1		2					25	765	790	26.88	22.96
(%)	(44.0)	(24.0)	(12.0)	(4.0)	(4.0)	(4.0)		(8.0)					(100.0)				
総計	11	67	2,396	8,953	11,008	7,231	2,910	1,073	371	109	46	31	34,206	765	34,971	56.35	12.88
(%)	(0.0)	(0.2)	(7.0)	(26.2)	(32.2)	(21.1)	(8.5)	(3.1)	(1.1)	(0.3)	(0.1)	(0.1)	(100.0)				
平均		1.61	1.88	2.18	2.51	2.83	3.20	3.62	4.02	4.38	4.80	5.19	2.57		2.57		
標準偏差		1.49	0.78	0.87	0.91	0.99	0.99	1.05	1.12	1.00	1.15	1.73	1.03		1.03		

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査