

表 2521 透析後体重(kg) 体重減少量(kg)別（血液透析濾過患者 男性のみ 透析回数週3回 透析歴2年以上）

体重減少量	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
0.0未満			2	8	23	16	7	2					58		58	59.22	10.32
(%)			(3.4)	(13.8)	(39.7)	(27.6)	(12.1)	(3.4)					(100.0)				
0.0～			26	173	250	150	47	11	3	1			661		661	56.24	10.34
(%)			(3.9)	(26.2)	(37.8)	(22.7)	(7.1)	(1.7)	(0.5)	(0.2)			(100.0)				
1.0～		2	103	876	1,448	789	178	38	6			1	3,441		3,441	55.57	9.42
(%)		(0.1)	(3.0)	(25.5)	(42.1)	(22.9)	(5.2)	(1.1)	(0.2)			(0.0)	(100.0)				
2.0～		2	98	1,509	3,568	2,332	727	165	40	6	2	1	8,450		8,450	58.27	9.75
(%)		(0.0)	(1.2)	(17.9)	(42.2)	(27.6)	(8.6)	(2.0)	(0.5)	(0.1)	(0.0)	(0.0)	(100.0)				
3.0～			22	502	1,969	2,172	1,047	357	116	30	4	4	6,223		6,223	63.67	11.30
(%)			(0.4)	(8.1)	(31.6)	(34.9)	(16.8)	(5.7)	(1.9)	(0.5)	(0.1)	(0.1)	(100.0)				
4.0～			1	55	374	584	492	297	113	37	15	10	1,978		1,978	71.11	13.86
(%)			(0.1)	(2.8)	(18.9)	(29.5)	(24.9)	(15.0)	(5.7)	(1.9)	(0.8)	(0.5)	(100.0)				
5.0～				4	28	83	88	65	57	28	15	7	375		375	80.46	17.02
(%)				(1.1)	(7.5)	(22.1)	(23.5)	(17.3)	(15.2)	(7.5)	(4.0)	(1.9)	(100.0)				
6.0～					8	6	6	8	13	3	4	4	52		52	87.27	23.76
(%)					(15.4)	(11.5)	(11.5)	(15.4)	(25.0)	(5.8)	(7.7)	(7.7)	(100.0)				
7.0～				1	3	2		1	1	1	2	2	13		13	86.16	31.53
(%)				(7.7)	(23.1)	(15.4)		(7.7)	(7.7)	(7.7)	(15.4)	(15.4)	(100.0)				
8.0～				5	11	2		1	1			2	22		22	63.57	25.47
(%)				(22.7)	(50.0)	(9.1)		(4.5)	(4.5)			(9.1)	(100.0)				
合計		4	252	3,133	7,682	6,136	2,592	945	350	106	42	31	21,273		21,273	61.03	12.14
(%)		(0.0)	(1.2)	(14.7)	(36.1)	(28.8)	(12.2)	(4.4)	(1.6)	(0.5)	(0.2)	(0.1)	(100.0)				
記載なし	8	2	2		1	1		2					16	487	503	29.46	27.27
(%)	(50.0)	(12.5)	(12.5)		(6.3)	(6.3)		(12.5)					(100.0)				
総計	8	6	254	3,133	7,683	6,137	2,592	947	350	106	42	31	21,289	487	21,776	61.01	12.19
(%)	(0.0)	(0.0)	(1.2)	(14.7)	(36.1)	(28.8)	(12.2)	(4.4)	(1.6)	(0.5)	(0.2)	(0.1)	(100.0)				
平均		1.83	1.88	2.28	2.58	2.87	3.23	3.65	4.05	4.41	4.90	5.19	2.78		2.78		
標準偏差		0.46	0.79	1.00	0.95	1.01	0.99	1.05	1.12	0.99	1.11	1.73	1.08		1.08		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査