

表 2522 透析後体重(kg) 体重減少量(kg)別（血液透析濾過患者 女性のみ 透析回数週3回 透析歴2年以上）

体重減少量	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
0.0未満		1	11	18	9								39		39	43.64	7.66
(%)		(2.6)	(28.2)	(46.2)	(23.1)								(100.0)				
0.0～		13	182	307	120	31	5	1					659		659	44.84	8.63
(%)		(2.0)	(27.6)	(46.6)	(18.2)	(4.7)	(0.8)	(0.2)					(100.0)				
1.0～		34	974	1,937	791	174	36	10	3				3,959		3,959	45.63	8.45
(%)		(0.9)	(24.6)	(48.9)	(20.0)	(4.4)	(0.9)	(0.3)	(0.1)				(100.0)				
2.0～		8	851	2,927	1,734	530	118	33	1	1	1		6,204		6,204	48.82	8.95
(%)		(0.1)	(13.7)	(47.2)	(27.9)	(8.5)	(1.9)	(0.5)	(0.0)	(0.0)	(0.0)		(100.0)				
3.0～			110	592	606	301	120	48	12	1	1		1,791		1,791	54.53	11.41
(%)			(6.1)	(33.1)	(33.8)	(16.8)	(6.7)	(2.7)	(0.7)	(0.1)	(0.1)		(100.0)				
4.0～			11	29	55	51	34	27	4	1	1		213		213	63.12	14.80
(%)			(5.2)	(13.6)	(25.8)	(23.9)	(16.0)	(12.7)	(1.9)	(0.5)	(0.5)		(100.0)				
5.0～			1	1	7	6	4	4	1		1		25		25	68.98	17.85
(%)			(4.0)	(4.0)	(28.0)	(24.0)	(16.0)	(16.0)	(4.0)		(4.0)		(100.0)				
6.0～					3	1		3					7		7	67.66	15.42
(%)					(42.9)	(14.3)		(42.9)					(100.0)				
7.0～				2			1						3		3	54.10	17.68
(%)				(66.7)			(33.3)						(100.0)				
8.0～		1	1	6									8		8	39.42	4.57
(%)		(12.5)	(12.5)	(75.0)									(100.0)				
合計		57	2,141	5,819	3,325	1,094	318	126	21	3	4		12,908		12,908	48.70	9.99
(%)		(0.4)	(16.6)	(45.1)	(25.8)	(8.5)	(2.5)	(1.0)	(0.2)	(0.0)	(0.0)		(100.0)				
記載なし	3	4	1	1									9	278	287	22.30	12.27
(%)	(33.3)	(44.4)	(11.1)	(11.1)									(100.0)				
総計	3	61	2,142	5,820	3,325	1,094	318	126	21	3	4		12,917	278	13,195	48.68	10.01
(%)	(0.0)	(0.5)	(16.6)	(45.1)	(25.7)	(8.5)	(2.5)	(1.0)	(0.2)	(0.0)	(0.0)		(100.0)				
平均		1.59	1.89	2.13	2.36	2.62	2.97	3.37	3.54	3.47	3.82		2.22		2.22		
標準偏差		1.54	0.78	0.79	0.78	0.85	0.89	1.05	1.09	1.04	1.26		0.84		0.84		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査