

表 2527 透析後体重(kg) Body Mass Index別（血液透析濾過患者 男性のみ 透析回数週3回）

Body Mass Index	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
14未満	2	4	30	6	1								43		43	35.64	7.41
(%)	(4.7)	(9.3)	(69.8)	(14.0)	(2.3)								(100.0)				
14～		1	127	229	4		1						362		362	41.50	4.10
(%)		(0.3)	(35.1)	(63.3)	(1.1)		(0.3)						(100.0)				
16～		1	82	1,310	453	1							1,847		1,847	46.90	4.26
(%)		(0.1)	(4.4)	(70.9)	(24.5)	(0.1)							(100.0)				
18～			16	1,388	2,765	206							4,375		4,375	52.09	4.62
(%)			(0.4)	(31.7)	(63.2)	(4.7)							(100.0)				
20～			3	366	3,524	1,678	51						5,622		5,622	57.51	5.03
(%)			(0.1)	(6.5)	(62.7)	(29.8)	(0.9)						(100.0)				
22～			1	28	1,302	2,862	478	5					4,676		4,676	63.05	5.47
(%)			(0.0)	(0.6)	(27.8)	(61.2)	(10.2)	(0.1)					(100.0)				
24～				4	183	1,528	1,092	101	1				2,909		2,909	68.82	6.05
(%)				(0.1)	(6.3)	(52.5)	(37.5)	(3.5)	(0.0)				(100.0)				
26～				1	10	341	865	329	22				1,568		1,568	75.09	6.71
(%)				(0.1)	(0.6)	(21.7)	(55.2)	(21.0)	(1.4)				(100.0)				
28～					2	47	313	358	91	4			815		815	81.06	7.27
(%)					(0.2)	(5.8)	(38.4)	(43.9)	(11.2)	(0.5)			(100.0)				
30～				1	4	9	72	247	251	79	10		673		673	90.04	9.39
(%)				(0.1)	(0.6)	(1.3)	(10.7)	(36.7)	(37.3)	(11.7)	(1.5)		(100.0)				
合計	2	6	259	3,333	8,248	6,672	2,872	1,040	365	83	10		22,890		22,890	60.89	11.54
(%)	(0.0)	(0.0)	(1.1)	(14.6)	(36.0)	(29.1)	(12.5)	(4.5)	(1.6)	(0.4)	(0.0)		(100.0)				
記載なし	7		31	248	521	434	175	87	63	54	47	37	1,704	600	2,304	65.39	19.74
(%)	(0.4)		(1.8)	(14.6)	(30.6)	(25.5)	(10.3)	(5.1)	(3.7)	(3.2)	(2.8)	(2.2)	(100.0)				
総計	9	6	290	3,581	8,769	7,106	3,047	1,127	428	137	57	37	24,594	600	25,194	61.21	12.34
(%)	(0.0)	(0.0)	(1.2)	(14.6)	(35.7)	(28.9)	(12.4)	(4.6)	(1.7)	(0.6)	(0.2)	(0.2)	(100.0)				
平均	6.08	12.63	15.73	18.19	20.53	23.07	25.77	28.57	31.01	32.87	33.23		22.11		22.11		
標準偏差	1.28	3.78	1.54	1.56	1.67	1.79	1.99	2.06	1.93	1.48	1.13		3.51		3.51		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査