

表 2528 透析後体重(kg) Body Mass Index別（血液透析濾過患者 女性のみ 透析回数週3回）

Body Mass Index	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
14未満	2	36	67	1									106		106	30.72	4.54
(%)	(1.9)	(34.0)	(63.2)	(0.9)									(100.0)				
14～		19	596	69									684		684	35.68	3.21
(%)		(2.8)	(87.1)	(10.1)									(100.0)				
16～		6	1,056	1,081	8								2,151		2,151	40.05	3.56
(%)		(0.3)	(49.1)	(50.3)	(0.4)								(100.0)				
18～		4	420	2,583	229	3							3,239		3,239	44.26	3.97
(%)		(0.1)	(13.0)	(79.7)	(7.1)	(0.1)							(100.0)				
20～			88	1,724	1,009	19							2,840		2,840	48.48	4.47
(%)			(3.1)	(60.7)	(35.5)	(0.7)							(100.0)				
22～		1	8	487	1,276	122	1						1,895		1,895	52.92	4.74
(%)		(0.1)	(0.4)	(25.7)	(67.3)	(6.4)	(0.1)						(100.0)				
24～			2	70	714	377	14	2					1,179		1,179	57.69	5.23
(%)			(0.2)	(5.9)	(60.6)	(32.0)	(1.2)	(0.2)					(100.0)				
26～				7	201	364	53	4					629		629	62.39	5.70
(%)				(1.1)	(32.0)	(57.9)	(8.4)	(0.6)					(100.0)				
28～				2	39	196	115	9	1				362		362	67.31	6.36
(%)				(0.6)	(10.8)	(54.1)	(31.8)	(2.5)	(0.3)				(100.0)				
30～				1	13	81	157	84	7	1			344		344	74.84	8.19
(%)				(0.3)	(3.8)	(23.5)	(45.6)	(24.4)	(2.0)	(0.3)			(100.0)				
合計	2	66	2,237	6,025	3,489	1,162	340	99	8	1			13,429		13,429	48.59	9.68
(%)	(0.0)	(0.5)	(16.7)	(44.9)	(26.0)	(8.7)	(2.5)	(0.7)	(0.1)	(0.0)			(100.0)				
記載なし	1	4	155	399	232	97	36	47	19	4	4		998	325	1,323	51.97	14.75
(%)	(0.1)	(0.4)	(15.5)	(40.0)	(23.2)	(9.7)	(3.6)	(4.7)	(1.9)	(0.4)	(0.4)		(100.0)				
総計	3	70	2,392	6,424	3,721	1,259	376	146	27	5	4		14,427	325	14,752	48.82	10.15
(%)	(0.0)	(0.5)	(16.6)	(44.5)	(25.8)	(8.7)	(2.6)	(1.0)	(0.2)	(0.0)	(0.0)		(100.0)				
平均	6.00	13.92	16.89	19.58	22.87	26.44	29.79	31.90	32.87	34.91			20.91		20.91		
標準偏差	1.32	2.52	1.65	1.81	2.09	2.27	2.16	2.05	1.42				3.75		3.75		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査