

表 2540 透析後体重(kg) %クレアチニン産生速度(%)別（血液透析濾過患者 透析回数週3回 透析歴2年以上）

%クレアチニン 産生速度	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
60未満		17	281	523	454	225	79	34	16	4	1		1,634		1,634	51.49	12.96
(%)		(1.0)	(17.2)	(32.0)	(27.8)	(13.8)	(4.8)	(2.1)	(1.0)	(0.2)	(0.1)		(100.0)				
60～		10	171	417	438	218	90	40	13	3	2	2	1,404		1,404	53.82	13.19
(%)		(0.7)	(12.2)	(29.7)	(31.2)	(15.5)	(6.4)	(2.8)	(0.9)	(0.2)	(0.1)	(0.1)	(100.0)				
70～		6	250	722	776	408	179	80	43	7	3	4	2,478		2,478	55.07	13.86
(%)		(0.2)	(10.1)	(29.1)	(31.3)	(16.5)	(7.2)	(3.2)	(1.7)	(0.3)	(0.1)	(0.2)	(100.0)				
80～		7	317	1,082	1,261	742	277	118	40	14	5	5	3,868		3,868	55.56	12.92
(%)		(0.2)	(8.2)	(28.0)	(32.6)	(19.2)	(7.2)	(3.1)	(1.0)	(0.4)	(0.1)	(0.1)	(100.0)				
90～		6	341	1,351	1,681	1,051	429	165	47	19	5	5	5,100		5,100	56.28	12.73
(%)		(0.1)	(6.7)	(26.5)	(33.0)	(20.6)	(8.4)	(3.2)	(0.9)	(0.4)	(0.1)	(0.1)	(100.0)				
100～		4	360	1,528	1,905	1,239	490	194	71	23	13	7	5,834		5,834	56.75	12.95
(%)		(0.1)	(6.2)	(26.2)	(32.7)	(21.2)	(8.4)	(3.3)	(1.2)	(0.4)	(0.2)	(0.1)	(100.0)				
110～			266	1,337	1,766	1,263	507	175	52	18	10	5	5,399		5,399	57.41	12.55
(%)			(4.9)	(24.8)	(32.7)	(23.4)	(9.4)	(3.2)	(1.0)	(0.3)	(0.2)	(0.1)	(100.0)				
120～		1	166	953	1,239	939	383	108	42	9	2	1	3,843		3,843	57.48	12.02
(%)		(0.0)	(4.3)	(24.8)	(32.2)	(24.4)	(10.0)	(2.8)	(1.1)	(0.2)	(0.1)	(0.0)	(100.0)				
130～			93	453	689	522	229	70	25	6	4	1	2,092		2,092	58.35	12.42
(%)			(4.4)	(21.7)	(32.9)	(25.0)	(10.9)	(3.3)	(1.2)	(0.3)	(0.2)	(0.0)	(100.0)				
140～		1	40	236	418	378	160	58	14	2		1	1,308		1,308	59.72	12.03
(%)		(0.1)	(3.1)	(18.0)	(32.0)	(28.9)	(12.2)	(4.4)	(1.1)	(0.2)		(0.1)	(100.0)				
合計		52	2,285	8,602	10,627	6,985	2,823	1,042	363	105	45	31	32,960		32,960	56.44	12.86
(%)		(0.2)	(6.9)	(26.1)	(32.2)	(21.2)	(8.6)	(3.2)	(1.1)	(0.3)	(0.1)	(0.1)	(100.0)				
記載なし	11	15	111	351	381	246	87	31	8	4	1		1,246	765	2,011	54.05	13.31
(%)	(0.9)	(1.2)	(8.9)	(28.2)	(30.6)	(19.7)	(7.0)	(2.5)	(0.6)	(0.3)	(0.1)		(100.0)				
総計	11	67	2,396	8,953	11,008	7,231	2,910	1,073	371	109	46	31	34,206	765	34,971	56.35	12.88
(%)	(0.0)	(0.2)	(7.0)	(26.2)	(32.2)	(21.1)	(8.5)	(3.1)	(1.1)	(0.3)	(0.1)	(0.1)	(100.0)				
平均		72.47	91.79	99.44	102.44	105.74	106.28	103.62	101.75	101.56	103.05	98.05	101.92		101.92		
標準偏差		25.12	26.83	23.74	23.20	23.16	22.95	23.07	22.96	21.45	18.44	19.44	23.87		23.87		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査