

表 2542 透析後体重(kg) %クレアチニン産生速度(%)別（血液透析濾過患者 女性のみ 透析回数週3回 透析歴2年以上）

%クレアチニン 産生速度	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
60未満		16	223	231	107	36	14	3	2				632		632	44.63	10.66
(%)		(2.5)	(35.3)	(36.6)	(16.9)	(5.7)	(2.2)	(0.5)	(0.3)				(100.0)				
60～		8	144	207	102	41	9	6	1				518		518	46.64	10.67
(%)		(1.5)	(27.8)	(40.0)	(19.7)	(7.9)	(1.7)	(1.2)	(0.2)				(100.0)				
70～		6	212	400	195	66	19	8	1				907		907	46.97	10.04
(%)		(0.7)	(23.4)	(44.1)	(21.5)	(7.3)	(2.1)	(0.9)	(0.1)				(100.0)				
80～		7	275	590	308	100	39	18	1	1			1,339		1,339	48.18	10.36
(%)		(0.5)	(20.5)	(44.1)	(23.0)	(7.5)	(2.9)	(1.3)	(0.1)	(0.1)			(100.0)				
90～		5	321	837	494	153	53	22	4	2	1		1,892		1,892	48.82	10.38
(%)		(0.3)	(17.0)	(44.2)	(26.1)	(8.1)	(2.8)	(1.2)	(0.2)	(0.1)	(0.1)		(100.0)				
100～		4	330	1,063	581	185	58	17	4				2,242		2,242	48.85	9.48
(%)		(0.2)	(14.7)	(47.4)	(25.9)	(8.3)	(2.6)	(0.8)	(0.2)				(100.0)				
110～			250	986	554	177	49	23	6		2		2,047		2,047	49.52	9.91
(%)			(12.2)	(48.2)	(27.1)	(8.6)	(2.4)	(1.1)	(0.3)		(0.1)		(100.0)				
120～		1	160	720	435	155	35	15	1				1,522		1,522	49.87	9.18
(%)		(0.1)	(10.5)	(47.3)	(28.6)	(10.2)	(2.3)	(1.0)	(0.1)				(100.0)				
130～			91	373	267	77	17	2	1		1		829		829	49.84	8.92
(%)			(11.0)	(45.0)	(32.2)	(9.3)	(2.1)	(0.2)	(0.1)		(0.1)		(100.0)				
140～		1	35	193	152	61	19	7					468		468	51.88	10.00
(%)		(0.2)	(7.5)	(41.2)	(32.5)	(13.0)	(4.1)	(1.5)					(100.0)				
合計		48	2,041	5,600	3,195	1,051	312	121	21	3	4		12,396		12,396	48.74	9.99
(%)		(0.4)	(16.5)	(45.2)	(25.8)	(8.5)	(2.5)	(1.0)	(0.2)	(0.0)	(0.0)		(100.0)				
記載なし	3	13	101	220	130	43	6	5					521	278	799	47.14	10.38
(%)	(0.6)	(2.5)	(19.4)	(42.2)	(25.0)	(8.3)	(1.2)	(1.0)					(100.0)				
総計	3	61	2,142	5,820	3,325	1,094	318	126	21	3	4		12,917	278	13,195	48.68	10.01
(%)	(0.0)	(0.5)	(16.6)	(45.1)	(25.7)	(8.5)	(2.5)	(1.0)	(0.2)	(0.0)	(0.0)		(100.0)				
平均		72.97	93.20	103.22	105.66	106.06	103.48	101.96	99.56	92.47	114.28		102.31		102.31		
標準偏差		25.52	25.86	22.74	22.83	24.31	23.06	21.84	21.03	3.13	15.27		23.89		23.89		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査