

表 2549 透析後体重(kg) 透析前ヘモグロビン濃度(g/dL)別（血液透析濾過患者 透析回数週3回）

透析前ヘモグロビン濃度	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
5未満			3	8	6	4	4	1					26		26	55.55	13.51
(%)			(11.5)	(30.8)	(23.1)	(15.4)	(15.4)	(3.8)					(100.0)				
5～			3	7	8	8	3	1	1				31		31	57.05	14.10
(%)			(9.7)	(22.6)	(25.8)	(25.8)	(9.7)	(3.2)	(3.2)				(100.0)				
6～		1	13	36	31	22	8	2					113	1	114	52.89	11.71
(%)		(0.9)	(11.5)	(31.9)	(27.4)	(19.5)	(7.1)	(1.8)					(100.0)				
7～		3	46	147	125	77	28	4	5		1		436	9	445	53.07	12.49
(%)		(0.7)	(10.6)	(33.7)	(28.7)	(17.7)	(6.4)	(0.9)	(1.1)		(0.2)		(100.0)				
8～	1	3	167	502	527	264	85	45	10	3	2	1	1,610	27	1,637	53.64	12.28
(%)	(0.1)	(0.2)	(10.4)	(31.2)	(32.7)	(16.4)	(5.3)	(2.8)	(0.6)	(0.2)	(0.1)	(0.1)	(100.0)				
9～	1	21	500	1,622	1,948	1,070	428	136	54	18	9	4	5,811	39	5,850	54.97	12.61
(%)	(0.0)	(0.4)	(8.6)	(27.9)	(33.5)	(18.4)	(7.4)	(2.3)	(0.9)	(0.3)	(0.2)	(0.1)	(100.0)				
10～	5	17	904	3,370	4,178	2,713	1,062	419	144	47	24	12	12,895	52	12,947	56.39	12.96
(%)	(0.0)	(0.1)	(7.0)	(26.1)	(32.4)	(21.0)	(8.2)	(3.2)	(1.1)	(0.4)	(0.2)	(0.1)	(100.0)				
11～	5	19	687	2,888	3,582	2,652	1,074	400	147	45	15	14	11,528	49	11,577	57.30	13.11
(%)	(0.0)	(0.2)	(6.0)	(25.1)	(31.1)	(23.0)	(9.3)	(3.5)	(1.3)	(0.4)	(0.1)	(0.1)	(100.0)				
12～		9	251	971	1,448	1,006	445	154	59	10	6	3	4,362	20	4,382	57.73	12.90
(%)		(0.2)	(5.8)	(22.3)	(33.2)	(23.1)	(10.2)	(3.5)	(1.4)	(0.2)	(0.1)	(0.1)	(100.0)				
13～			55	210	330	313	174	55	21	11	4	2	1,175	10	1,185	60.73	13.97
(%)			(4.7)	(17.9)	(28.1)	(26.6)	(14.8)	(4.7)	(1.8)	(0.9)	(0.3)	(0.2)	(100.0)				
14～			16	93	138	131	83	39	9	5		1	515	8	523	62.01	14.06
(%)			(3.1)	(18.1)	(26.8)	(25.4)	(16.1)	(7.6)	(1.7)	(1.0)		(0.2)	(100.0)				
合計	12	73	2,645	9,854	12,321	8,260	3,394	1,256	450	139	61	37	38,502	215	38,717	56.64	13.04
(%)	(0.0)	(0.2)	(6.9)	(25.6)	(32.0)	(21.5)	(8.8)	(3.3)	(1.2)	(0.4)	(0.2)	(0.1)	(100.0)				
記載なし		3	37	151	169	105	29	17	5	3			519	710	1,229	55.32	12.53
(%)		(0.6)	(7.1)	(29.1)	(32.6)	(20.2)	(5.6)	(3.3)	(1.0)	(0.6)			(100.0)				
総計	12	76	2,682	10,005	12,490	8,365	3,423	1,273	455	142	61	37	39,021	925	39,946	56.63	13.03
(%)	(0.0)	(0.2)	(6.9)	(25.6)	(32.0)	(21.4)	(8.8)	(3.3)	(1.2)	(0.4)	(0.2)	(0.1)	(100.0)				
平均	10.67	10.38	10.59	10.71	10.82	10.96	11.07	11.08	11.06	11.18	10.79	11.12	10.84	10.60	10.84		
標準偏差	0.92	1.32	1.28	1.24	1.23	1.25	1.32	1.30	1.27	1.37	1.19	1.34	1.26	1.64	1.26		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査