

表 2551 透析後体重(kg) HDL-C濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

HDL-C濃度	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
20未満			4	18	30	14	7	2	4	2			81		81	59.58	15.83
(%)			(4.9)	(22.2)	(37.0)	(17.3)	(8.6)	(2.5)	(4.9)	(2.5)			(100.0)				
20～			55	255	449	486	320	168	81	27	16	10	1,867	7	1,874	65.13	15.99
(%)			(2.9)	(13.7)	(24.0)	(26.0)	(17.1)	(9.0)	(4.3)	(1.4)	(0.9)	(0.5)	(100.0)				
30～	2	7	226	1,006	1,767	1,665	813	373	163	47	15	14	6,098	33	6,131	61.51	14.27
(%)	(0.0)	(0.1)	(3.7)	(16.5)	(29.0)	(27.3)	(13.3)	(6.1)	(2.7)	(0.8)	(0.2)	(0.2)	(100.0)				
40～	2	11	392	1,652	2,446	1,733	737	250	69	19	9	7	7,327	33	7,360	57.61	12.56
(%)	(0.0)	(0.2)	(5.4)	(22.5)	(33.4)	(23.7)	(10.1)	(3.4)	(0.9)	(0.3)	(0.1)	(0.1)	(100.0)				
50～		15	441	1,652	2,073	1,218	395	113	34	5	5	1	5,952	19	5,971	54.86	11.41
(%)		(0.3)	(7.4)	(27.8)	(34.8)	(20.5)	(6.6)	(1.9)	(0.6)	(0.1)	(0.1)	(0.0)	(100.0)				
60～	1	9	340	1,351	1,319	605	166	44	7	3	2		3,847	10	3,857	52.57	10.49
(%)	(0.0)	(0.2)	(8.8)	(35.1)	(34.3)	(15.7)	(4.3)	(1.1)	(0.2)	(0.1)	(0.1)		(100.0)				
70～	2	9	236	721	647	295	73	27	6	3			2,019	4	2,023	51.61	10.75
(%)	(0.1)	(0.4)	(11.7)	(35.7)	(32.0)	(14.6)	(3.6)	(1.3)	(0.3)	(0.1)			(100.0)				
80～		3	118	399	324	131	30	5	2				1,012	1	1,013	50.79	9.78
(%)		(0.3)	(11.7)	(39.4)	(32.0)	(12.9)	(3.0)	(0.5)	(0.2)				(100.0)				
90～			59	169	134	57	9	2					430	1	431	49.97	9.79
(%)			(13.7)	(39.3)	(31.2)	(13.3)	(2.1)	(0.5)					(100.0)				
100～			64	134	112	59	9	2	1				381	2	383	50.24	10.14
(%)			(16.8)	(35.2)	(29.4)	(15.5)	(2.4)	(0.5)	(0.3)				(100.0)				
合計	7	54	1,935	7,357	9,301	6,263	2,559	986	367	106	47	32	29,014	110	29,124	56.82	13.12
(%)	(0.0)	(0.2)	(6.7)	(25.4)	(32.1)	(21.6)	(8.8)	(3.4)	(1.3)	(0.4)	(0.2)	(0.1)	(100.0)				
記載なし	5	22	747	2,648	3,189	2,102	864	287	88	36	14	5	10,007	815	10,822	56.06	12.77
(%)	(0.0)	(0.2)	(7.5)	(26.5)	(31.9)	(21.0)	(8.6)	(2.9)	(0.9)	(0.4)	(0.1)	(0.0)	(100.0)				
総計	12	76	2,682	10,005	12,490	8,365	3,423	1,273	455	142	61	37	39,021	925	39,946	56.63	13.03
(%)	(0.0)	(0.2)	(6.9)	(25.6)	(32.0)	(21.4)	(8.8)	(3.3)	(1.2)	(0.4)	(0.2)	(0.1)	(100.0)				
平均	54.43	56.17	58.26	55.79	51.68	47.51	43.34	40.49	37.27	36.25	35.81	33.75	50.87	46.96	50.85		
標準偏差	17.21	14.56	19.00	17.45	16.67	15.91	13.85	12.79	11.83	11.49	11.05	8.02	17.18	16.24	17.18		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査