

表 2552 透析後体重(kg) Non-HDL-C濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

Non-HDL-C濃度	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
50未満		1	37	127	190	107	48	10	5	2			527	2	529	55.83	11.88
(%)		(0.2)	(7.0)	(24.1)	(36.1)	(20.3)	(9.1)	(1.9)	(0.9)	(0.4)			(100.0)				
50～	3	2	137	577	842	559	179	67	21	7	2	1	2,397	6	2,403	56.61	12.04
(%)	(0.1)	(0.1)	(5.7)	(24.1)	(35.1)	(23.3)	(7.5)	(2.8)	(0.9)	(0.3)	(0.1)	(0.0)	(100.0)				
70～	2	8	320	1,294	1,776	1,105	408	143	45	11	6	1	5,119	19	5,138	56.20	12.09
(%)	(0.0)	(0.2)	(6.3)	(25.3)	(34.7)	(21.6)	(8.0)	(2.8)	(0.9)	(0.2)	(0.1)	(0.0)	(100.0)				
90～	1	10	426	1,634	2,050	1,330	516	169	70	17	10	3	6,236	29	6,265	56.25	12.58
(%)	(0.0)	(0.2)	(6.8)	(26.2)	(32.9)	(21.3)	(8.3)	(2.7)	(1.1)	(0.3)	(0.2)	(0.0)	(100.0)				
110～		11	370	1,319	1,570	1,098	482	204	58	20	11	6	5,149	13	5,162	56.96	13.49
(%)		(0.2)	(7.2)	(25.6)	(30.5)	(21.3)	(9.4)	(4.0)	(1.1)	(0.4)	(0.2)	(0.1)	(100.0)				
130～		4	221	808	914	668	306	127	62	15	3	10	3,138	11	3,149	57.61	14.17
(%)		(0.1)	(7.0)	(25.7)	(29.1)	(21.3)	(9.8)	(4.0)	(2.0)	(0.5)	(0.1)	(0.3)	(100.0)				
150～		3	104	374	474	344	182	61	33	9	5	4	1,593	4	1,597	58.22	14.40
(%)		(0.2)	(6.5)	(23.5)	(29.8)	(21.6)	(11.4)	(3.8)	(2.1)	(0.6)	(0.3)	(0.3)	(100.0)				
170～	1		48	172	196	136	67	38	14	8	1	1	682		682	58.19	14.96
(%)	(0.1)		(7.0)	(25.2)	(28.7)	(19.9)	(9.8)	(5.6)	(2.1)	(1.2)	(0.1)	(0.1)	(100.0)				
190～		1	16	74	64	57	26	17	4	2		1	262		262	57.94	14.93
(%)		(0.4)	(6.1)	(28.2)	(24.4)	(21.8)	(9.9)	(6.5)	(1.5)	(0.8)		(0.4)	(100.0)				
210～			8	36	45	25	15	6	5	1			141		141	58.26	14.41
(%)			(5.7)	(25.5)	(31.9)	(17.7)	(10.6)	(4.3)	(3.5)	(0.7)			(100.0)				
合計	7	40	1,687	6,415	8,121	5,429	2,229	842	317	92	38	27	25,244	84	25,328	56.78	13.06
(%)	(0.0)	(0.2)	(6.7)	(25.4)	(32.2)	(21.5)	(8.8)	(3.3)	(1.3)	(0.4)	(0.2)	(0.1)	(100.0)				
記載なし	5	36	995	3,590	4,369	2,936	1,194	431	138	50	23	10	13,777	841	14,618	56.34	12.97
(%)	(0.0)	(0.3)	(7.2)	(26.1)	(31.7)	(21.3)	(8.7)	(3.1)	(1.0)	(0.4)	(0.2)	(0.1)	(100.0)				
総計	12	76	2,682	10,005	12,490	8,365	3,423	1,273	455	142	61	37	39,021	925	39,946	56.63	13.03
(%)	(0.0)	(0.2)	(6.9)	(25.6)	(32.0)	(21.4)	(8.8)	(3.3)	(1.2)	(0.4)	(0.2)	(0.1)	(100.0)				
平均	90.43	109.10	108.33	107.35	104.91	106.50	110.28	113.90	117.96	120.33	114.21	131.00	107.14	101.44	107.12		
標準偏差	40.96	31.62	32.67	33.14	33.06	33.11	34.03	35.20	36.97	37.71	29.22	30.26	33.39	27.80	33.37		

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査