

表 2555 透析後体重(kg) 透析前脈拍(拍/分)別（血液透析濾過患者 透析回数週3回）

透析前脈拍	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
40未満			1	5	8	2	1						17		17	52.84	10.39
(%)			(5.9)	(29.4)	(47.1)	(11.8)	(5.9)						(100.0)				
40～		1	25	88	96	64	13	8					295		295	53.80	10.97
(%)		(0.3)	(8.5)	(29.8)	(32.5)	(21.7)	(4.4)	(2.7)					(100.0)				
50～	1	3	225	755	942	518	139	38	5	1	1		2,628	4	2,632	53.87	10.73
(%)	(0.0)	(0.1)	(8.6)	(28.7)	(35.8)	(19.7)	(5.3)	(1.4)	(0.2)	(0.0)	(0.0)		(100.0)				
60～	4	19	612	2,419	2,888	1,771	551	164	40	9	2	2	8,481	9	8,490	54.80	11.41
(%)	(0.0)	(0.2)	(7.2)	(28.5)	(34.1)	(20.9)	(6.5)	(1.9)	(0.5)	(0.1)	(0.0)	(0.0)	(100.0)				
70～	2	20	753	2,752	3,375	2,282	877	315	96	37	12	2	10,523	22	10,545	56.19	12.57
(%)	(0.0)	(0.2)	(7.2)	(26.2)	(32.1)	(21.7)	(8.3)	(3.0)	(0.9)	(0.4)	(0.1)	(0.0)	(100.0)				
80～	1	11	416	1,696	2,150	1,525	768	288	134	39	20	9	7,057	6	7,063	58.15	13.98
(%)	(0.0)	(0.2)	(5.9)	(24.0)	(30.5)	(21.6)	(10.9)	(4.1)	(1.9)	(0.6)	(0.3)	(0.1)	(100.0)				
90～	1	7	170	655	974	746	413	197	84	22	13	8	3,290	3	3,293	60.18	14.88
(%)	(0.0)	(0.2)	(5.2)	(19.9)	(29.6)	(22.7)	(12.6)	(6.0)	(2.6)	(0.7)	(0.4)	(0.2)	(100.0)				
100～		5	74	194	274	217	135	82	31	12	7	10	1,041	1	1,042	61.49	17.30
(%)		(0.5)	(7.1)	(18.6)	(26.3)	(20.8)	(13.0)	(7.9)	(3.0)	(1.2)	(0.7)	(1.0)	(100.0)				
110～		1	21	54	79	56	22	11	9	4		3	260	1	261	59.21	16.87
(%)		(0.4)	(8.1)	(20.8)	(30.4)	(21.5)	(8.5)	(4.2)	(3.5)	(1.5)		(1.2)	(100.0)				
120～			10	25	26	22	16	5	2				106		106	57.97	14.04
(%)			(9.4)	(23.6)	(24.5)	(20.8)	(15.1)	(4.7)	(1.9)				(100.0)				
合計	9	67	2,307	8,643	10,812	7,203	2,935	1,108	401	124	55	34	33,698	46	33,744	56.63	13.07
(%)	(0.0)	(0.2)	(6.8)	(25.6)	(32.1)	(21.4)	(8.7)	(3.3)	(1.2)	(0.4)	(0.2)	(0.1)	(100.0)				
記載なし	3	9	375	1,362	1,678	1,162	488	165	54	18	6	3	5,323	879	6,202	56.60	12.77
(%)	(0.1)	(0.2)	(7.0)	(25.6)	(31.5)	(21.8)	(9.2)	(3.1)	(1.0)	(0.3)	(0.1)	(0.1)	(100.0)				
総計	12	76	2,682	10,005	12,490	8,365	3,423	1,273	455	142	61	37	39,021	925	39,946	56.63	13.03
(%)	(0.0)	(0.2)	(6.9)	(25.6)	(32.0)	(21.4)	(8.8)	(3.3)	(1.2)	(0.4)	(0.2)	(0.1)	(100.0)				
平均	69.33	76.27	74.26	73.96	74.61	75.78	78.78	80.97	84.14	84.26	85.98	93.03	75.43	73.98	75.43		
標準偏差	11.54	14.33	13.27	12.49	12.72	12.81	13.02	13.32	12.03	11.81	11.42	12.81	12.93	12.74	12.93		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査