

表 2559 体重(kg) 透析歴(年)別（腹膜透析患者）

透析歴	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
2年未満	22	4	86	343	606	649	341	122	50	4	7	4	2,238	1,362	3,600	60.88	14.54
(%)	(1.0)	(0.2)	(3.8)	(15.3)	(27.1)	(29.0)	(15.2)	(5.5)	(2.2)	(0.2)	(0.3)	(0.2)	(100.0)				
2年～	11	10	44	271	568	585	299	120	35	9	1	3	1,956	1,245	3,201	61.69	13.41
(%)	(0.6)	(0.5)	(2.2)	(13.9)	(29.0)	(29.9)	(15.3)	(6.1)	(1.8)	(0.5)	(0.1)	(0.2)	(100.0)				
5年～	4	7	33	212	310	291	144	46	18	7	2	2	1,076	581	1,657	59.79	13.97
(%)	(0.4)	(0.7)	(3.1)	(19.7)	(28.8)	(27.0)	(13.4)	(4.3)	(1.7)	(0.7)	(0.2)	(0.2)	(100.0)				
10年～	1		3	50	75	64	19	10					222	118	340	58.31	10.81
(%)	(0.5)		(1.4)	(22.5)	(33.8)	(28.8)	(8.6)	(4.5)					(100.0)				
15年～		1	5	16	23	20	6		1				72	25	97	56.04	11.11
(%)		(1.4)	(6.9)	(22.2)	(31.9)	(27.8)	(8.3)		(1.4)				(100.0)				
20年～		1		5	4	4							14	6	20	52.01	10.47
(%)		(7.1)		(35.7)	(28.6)	(28.6)							(100.0)				
25年～				3	1	2	1						7	1	8	56.01	12.28
(%)				(42.9)	(14.3)	(28.6)	(14.3)						(100.0)				
30年～				3	4								7	3	10	48.37	6.66
(%)				(42.9)	(57.1)								(100.0)				
35年～			1	1									2	1	3	43.15	8.27
(%)			(50.0)	(50.0)									(100.0)				
40年～																	
(%)																	
合計	38	23	172	904	1,591	1,615	810	298	104	20	10	9	5,594	3,342	8,936	60.74	13.90
(%)	(0.7)	(0.4)	(3.1)	(16.2)	(28.4)	(28.9)	(14.5)	(5.3)	(1.9)	(0.4)	(0.2)	(0.2)	(100.0)				
不明						1	1						2	3	5	69.25	12.09
(%)						(50.0)	(50.0)						(100.0)				
記載なし																	
(%)																	
総計	38	23	172	904	1,591	1,616	811	298	104	20	10	9	5,596	3,345	8,941	60.74	13.90
(%)	(0.7)	(0.4)	(3.1)	(16.2)	(28.4)	(28.9)	(14.5)	(5.3)	(1.9)	(0.4)	(0.2)	(0.2)	(100.0)				
平均	2.24	4.91	3.06	3.71	3.37	3.08	2.76	2.63	2.32	3.40	2.00	2.11	3.18	2.97	3.10		
標準偏差	2.58	5.02	4.44	4.46	3.79	3.50	3.00	2.67	2.59	1.98	3.06	2.47	3.68	3.36	3.56		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査