

表 2572 体重(kg) 腎移植の回数別（腹膜透析患者）

腎移植の回数	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
なし	36	21	151	803	1,399	1,404	712	260	88	19	8	8	4,909	866	5,775	60.63	13.97
(%)	(0.7)	(0.4)	(3.1)	(16.4)	(28.5)	(28.6)	(14.5)	(5.3)	(1.8)	(0.4)	(0.2)	(0.2)	(100.0)				
1回	1			11	14	14	2	1				1	44	12	56	58.46	15.32
(%)	(2.3)			(25.0)	(31.8)	(31.8)	(4.5)	(2.3)				(2.3)	(100.0)				
2回	1												1	1	2	19.80	
(%)	(100.0)												(100.0)				
3回														1	1		
(%)																	
4回以上																	
(%)																	
合計	38	21	151	814	1,413	1,418	714	261	88	19	8	9	4,954	880	5,834	60.60	14.00
(%)	(0.8)	(0.4)	(3.0)	(16.4)	(28.5)	(28.6)	(14.4)	(5.3)	(1.8)	(0.4)	(0.2)	(0.2)	(100.0)				
不明				1	4	6							11	12	23	62.04	7.41
(%)				(9.1)	(36.4)	(54.5)							(100.0)				
記載なし		2	21	89	174	192	97	37	16	1	2		631	2,453	3,084	61.83	13.14
(%)		(0.3)	(3.3)	(14.1)	(27.6)	(30.4)	(15.4)	(5.9)	(2.5)	(0.2)	(0.3)		(100.0)				
総計	38	23	172	904	1,591	1,616	811	298	104	20	10	9	5,596	3,345	8,941	60.74	13.90
(%)	(0.7)	(0.4)	(3.1)	(16.2)	(28.4)	(28.9)	(14.5)	(5.3)	(1.9)	(0.4)	(0.2)	(0.2)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査