

表 2578 体重(kg) Body Mass Index別（腹膜透析患者 女性のみ）

Body Mass Index	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
14未満	7	4	4										15		15	21.15	10.88
(%)	(46.7)	(26.7)	(26.7)										(100.0)				
14～	10	5	28	5									48		48	29.99	11.17
(%)	(20.8)	(10.4)	(58.3)	(10.4)									(100.0)				
16～	2	3	60	76	2								143		143	39.84	5.28
(%)	(1.4)	(2.1)	(42.0)	(53.1)	(1.4)								(100.0)				
18～	1	1	42	262	30								336		336	44.49	4.57
(%)	(0.3)	(0.3)	(12.5)	(78.0)	(8.9)								(100.0)				
20～			12	226	189	3							430		430	49.05	4.59
(%)			(2.8)	(52.6)	(44.0)	(0.7)							(100.0)				
22～		1		96	240	29							366		366	53.17	5.17
(%)		(0.3)		(26.2)	(65.6)	(7.9)							(100.0)				
24～				10	140	83	4	1					238		238	58.05	5.29
(%)				(4.2)	(58.8)	(34.9)	(1.7)	(0.4)					(100.0)				
26～			1	1	45	92	15	2					156		156	62.87	6.50
(%)			(0.6)	(0.6)	(28.8)	(59.0)	(9.6)	(1.3)					(100.0)				
28～				2	11	55	31	4	1				104		104	67.85	7.54
(%)				(1.9)	(10.6)	(52.9)	(29.8)	(3.8)	(1.0)				(100.0)				
30～					2	17	37	15					71		71	73.70	7.00
(%)					(2.8)	(23.9)	(52.1)	(21.1)					(100.0)				
合計	20	14	147	678	659	279	87	22	1				1,907		1,907	51.84	10.94
(%)	(1.0)	(0.7)	(7.7)	(35.6)	(34.6)	(14.6)	(4.6)	(1.2)	(0.1)				(100.0)				
記載なし			7	22	31	8	8	3	9	1		3	92	1,200	1,292	62.32	23.54
(%)			(7.6)	(23.9)	(33.7)	(8.7)	(8.7)	(3.3)	(9.8)	(1.1)		(3.3)	(100.0)				
総計	20	14	154	700	690	287	95	25	10	1		3	1,999	1,200	3,199	52.33	12.01
(%)	(1.0)	(0.7)	(7.7)	(35.0)	(34.5)	(14.4)	(4.8)	(1.3)	(0.5)	(0.1)		(0.2)	(100.0)				
平均	14.44	15.41	17.44	20.11	23.09	26.56	29.56	31.26	29.90				22.35		22.35		
標準偏差	1.95	2.86	1.86	1.89	2.11	2.24	1.99	2.65					3.86		3.86		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査